

REVIEW

Convergent pharmacological mechanisms in impulsivity and addiction: insights from rodent models

B Jupp^{1,2} and J W Dalley^{1,3}

¹Behavioural and Clinical Neuroscience Institute and Department of Psychology, University of Cambridge, Cambridge, UK, ²Florey Institute of Neuroscience and Mental Health, University of Melbourne, Parkville, Australia, and ³Department of Psychiatry, University of Cambridge, Cambridge, UK

Correspondence

Jeffrey W Dalley, Department of Psychology, University of Cambridge, Downing St, Cambridge CB2 3EB, UK. E-mail: jwd20@cam.ac.uk

Received

20 February 2014

Revised

2 May 2014

Accepted

12 May 2014

Research over the last two decades has widely demonstrated that impulsivity, in its various forms, is antecedent to the development of drug addiction and an important behavioural trait underlying the inability of addicts to refrain from continued drug use. Impulsivity describes a variety of rapidly and prematurely expressed behaviours that span several domains from impaired response inhibition to an intolerance of delayed rewards, and is a core symptom of attention deficit hyperactivity disorder (ADHD) and other brain disorders. Various theories have been advanced to explain how impulsivity interacts with addiction both causally and as a consequence of chronic drug abuse; these acknowledge the strong overlaps in neural circuitry and mechanisms between impulsivity and addiction and the seemingly paradoxical treatment of ADHD with stimulant drugs with high abuse potential. Recent years have witnessed unprecedented progress in the elucidation of pharmacological mechanisms underpinning impulsivity. Collectively, this work has significantly improved the prospect for new therapies in ADHD as well as our understanding of the neural mechanisms underlying the shift from recreational drug use to addiction. In this review, we consider the extent to which pharmacological interventions that target impulsive behaviour are also effective in animal models of addiction. We highlight several promising examples of convergence based on empirical findings in rodent-based studies.

LINKED ARTICLES

This article is part of a themed section on Animal Models in Psychiatry Research. To view the other articles in this section visit <http://dx.doi.org/10.1111/bph.2014.171.issue-20>

Abbreviations

5-CSRTT, five choice serial reaction time task; ADHD, attention deficit hyperactivity disorder; DD, delayed discounting; DRL, differential reinforcement of lower rates; FR, fixed ratio; HI, highly impulsive; ILC, infralimbic cortex; NAc, nucleus accumbens; NARI, noradrenaline re-uptake inhibitor; OFC, orbitofrontal cortex; PFC, prefrontal cortex; PR, progressive ratio; SSRI, selective 5-HT re-uptake inhibitor; SSRTT, stop-signal reaction time task; STN, subthalamic nucleus; vHC, ventral hippocampus; VTA, ventral tegmental area

Table of Links

TARGETS	LIGANDS
α_1 -adrenoceptor	Dopamine
α_2 -adrenoceptor	5-HT
β_1 -adrenoceptor	SB242084
β_2 -adrenoceptor	WAY163909
D ₁ receptor	Ketanserin
D ₂ receptor	Desipramine
D ₃ receptor	Yohimbine
5-HT ₁ receptor	MK-801 (dizocilpine)
5-HT ₂ receptor	GABA
5-HT ₃ receptor	SR141716A (rimonabant)
5-HT ₆ receptor	Neuropeptide Y
GluN2B receptor	Methamphetamine
mGlu ₁ receptor	Cocaine
mGlu ₂ receptor	Ethanol
mGlu ₅ receptor	Atomoxetine
mGlu ₇ receptor	Amphetamine
GABA _A receptor	
GABA _B receptor	
μ -opioid receptor	
δ -opioid receptor	
CB ₁ receptor	
Nicotinic $\alpha 4\beta 2$ receptor	
M ₁ receptor	
M ₄ receptor	
NK ₁ receptor	
Y ₂ receptor	
H ₃ receptor	
A ₁ receptor	

This Table lists key protein targets and ligands in this document, which are hyperlinked to corresponding entries in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS Guide to PHARMACOLOGY (Pawson *et al.*, 2014) and are permanently archived in the Concise Guide to PHARMACOLOGY 2013/14 (Alexander *et al.*, 2013a, Alexander *et al.*, 2013b).

Introduction

Drug addiction is a chronic, relapsing brain disorder for which surprisingly few effective therapies have been developed (O'Brien, 2008; Jupp and Lawrence, 2010; van den Brink, 2012; Pierce *et al.*, 2012). Although the precise brain mechanisms of addiction are unknown, this disorder is widely believed to engage individual biological risk factors that interact individually and collectively with drug-, stress- and other externally influenced brain plasticity mechanisms (Nestler, 2005; Uhl, 2006; Koob, 2008; Kalivas and Volkow, 2011). Consistent with this interpretation, personality traits such as novelty/sensation-seeking and impulsivity are widely recognized to predispose to addiction and modify disease progression (Chakroun *et al.*, 2004; Nigg *et al.*, 2006; Verdejo-Garcia *et al.*, 2008; Ersche *et al.*, 2010). Such traits

share strong overlaps with structural and functional markers of addiction (Jupp and Dalley, 2014) and may, as a result, inform the neurobiological and pharmacological mechanisms of this disorder, thereby raising the possibility that modulating impulsivity may provide an approach through which risk for addiction and/or the addiction process may be remediated. Notably, in this regard, stimulant drugs such as d-amphetamine, which calm behaviour and reduce hyperactivity and impulsivity in attention deficit hyperactivity disorder (ADHD; Solanto, 1984; Fone and Nutt, 2005) are also prominent drugs of abuse. As a consequence, it has become an important open question whether treating ADHD with stimulant drugs accelerates, or conversely, offers protection against, the development of addiction (e.g. Barkley *et al.*, 2003). This paper reviews the evidence for convergent pharmacological mechanisms in impulsivity and addiction,

and additionally considers whether stimulant- and non-stimulant-based medications in ADHD may modify an individual's risk for addiction.

Defining impulsivity

Impulsivity describes an individual's tendency for premature, excessively risky, poorly conceived and inappropriate actions without due regard for future consequences (Daruma and Barnes, 1993). A range of behavioural processes are generally considered to contribute to this trait including urgency, risk-taking, sensation-seeking, behavioural disinhibition, impaired planning, lack of premeditation and insensitivity to punishment (Barratt, 1985; Evenden, 1999a; Monterosso and Ainslie, 1999; Moeller *et al.*, 2001; Whiteside and Lynam, 2003; Robbins *et al.*, 2012; Fineberg *et al.*, 2014). While impulsiveness is an important aspect of normal human behaviour, facilitating extraversion, sociability and appropriate risk-taking, the maladaptive expression of this trait has been associated with a number of neuropsychiatric morbidities including personality (Perry and Korner, 2011) and mood disorders (Lombardo *et al.*, 2012), suicide (Dougherty *et al.*, 2004), ADHD (Avila *et al.*, 2004), problem gambling (Verdejo-Garcia *et al.*, 2008) and drug addiction (Ersche *et al.*, 2010).

Impulsivity is a multifactorial trait often segregated according to motor disinhibition ('impulsive action') and impulsive decision making ('impulsive choice') (Winstanley *et al.*, 2006). In humans, it is generally assessed by self-report scales; for example, the Barratt Impulsivity Scale (BIS-11) (Barratt, 1985), the Urgency, Premeditation, Perseverance, Sensation-Seeking (UPSS) Impulsive Behaviour Scale (Whiteside and Lynam, 2003) and Dickman's Impulsivity Inventory (DII) (Dickman, 1990). Psychometric laboratory-based tasks can also be used to assess impulsivity; these overcome many of the limitations associated with self-report (e.g. see Wilson and Dunn, 2004) by providing more objective behavioural measures (Kertzman *et al.*, 2006; Chamberlain and Sahakian, 2007). In experimental animals, impulsive choice is frequently measured by operant-based delay-discounting tasks (Reynolds *et al.*, 2002; Cardinal, 2006). While several variants of this task are used, the general procedure involves a choice between a small immediate reward and a larger, but delayed reward. Impulsive choice is indexed by steeper reward discounting such that with increasing delay the perceived 'value' of reward diminishes and preference switches to small, immediate rewards. Action impulsivity can be assessed using a variety of operant paradigms; for example go/no-go visual and spatial discrimination tasks (Harrison *et al.*, 1999), which require inhibition of incorrect responses; the stop-signal reaction time task (SSRTT) where already initiated actions must be rapidly cancelled following presentation of an auditory or visual 'stop signal' stimulus (Eagle and Robbins, 2003). Impulsivity is assessed in these tasks by the number of inappropriate responses made. These include premature responses to a food-predictive cue on the five choice serial reaction time task (5-CSRTT) (Robbins, 2002), analogous to the differential responding for differential reinforcement of lower rate (DRL) task where responses made before a

time interval has elapsed are punished (Evenden and Ryan, 1996; Evenden, 1999a).

Impulsivity and addiction

Impulsivity is widely regarded to contribute to the development of addiction, predicting initial drug use (Zernicke *et al.*, 2010), risk for addiction (Ersche *et al.*, 2010), rates of relapse (Muller *et al.*, 2008) and treatment retention (Moeller *et al.*, 2001). However, drugs of abuse can in turn affect levels of impulsivity (Jentsch and Taylor, 1999; Garavan *et al.*, 2008; de Wit, 2009) making it unclear whether co-expressed impulsivity in addicts (e.g. Petry, 2001; Moreno-Lopez *et al.*, 2012) is a cause or consequence of chronic drug use (Rogers and Robbins, 2001). In contrast, animal models of impulsivity help disambiguate causal relationships between impulsivity and addiction by enabling assessment prior to, and following, chronic drug exposure (Winstanley *et al.*, 2010a; Jupp *et al.*, 2013a). Collectively, these studies demonstrate that impulsivity affects different measures of addiction-related behaviour depending on drug class, baseline levels of impulsivity and specific impulsivity subtypes; they further demonstrate that drug use can, in turn, affect levels of impulsivity. For example, action impulsivity in rats precedes enhanced self-administration of a range of drugs, including stimulants (Dalley *et al.*, 2007; Belin *et al.*, 2008; Marusich and Bardo, 2009), alcohol (Radwanska and Kaczmarek, 2012) and nicotine (Diergaarde *et al.*, 2008), and predicts increased rates of relapse to cocaine-seeking (Economidou *et al.*, 2009). Similarly, choice impulsivity predicts increased alcohol (Poulos *et al.*, 1995; Oberlin and Grahame, 2009) and nicotine (Diergaarde *et al.*, 2008; Kayir *et al.*, 2014) consumption in rats, as well as resistance to extinction and enhanced relapse propensity to both nicotine (Diergaarde *et al.*, 2008) and cocaine (Broos *et al.*, 2012a). However, the relationship between choice impulsivity and drug reinforcement is not always clear (Broos *et al.*, 2012a; Schippers *et al.*, 2012), and this may reflect direct modulatory effects of cocaine and heroin on this particular measure of impulsivity (Mendez *et al.*, 2010; Schippers *et al.*, 2012).

Intriguingly, the effect of certain drugs on impulsivity often depends on baseline levels of impulsivity. For example, cocaine (Paine *et al.*, 2003; Roesch *et al.*, 2007; Winstanley *et al.*, 2009; Mendez *et al.*, 2010; Caprioli *et al.*, 2013) and nicotine (Kayir *et al.*, 2014) are reported to increase impulsivity in non-impulsive rats, but have the opposite effect in impulsive animals (Dalley *et al.*, 2007; Caprioli *et al.*, 2013; Kayir *et al.*, 2014; Kolokotroni *et al.*, 2014). It follows therefore that enhanced drug intake in impulsive animals may represent a form of 'self-medication' (Khantzian, 1985) analogous to the treatment of ADHD with stimulant drugs. Furthermore, the close interrelationship between impulsivity and addiction implies that they may share similar psychobiological mechanisms (e.g. an intolerance of delayed rewards) and that interventions that reduce impulsivity would have clinical benefits in addiction. In this review, we consider the strength of evidence implicating shared pharmacological mechanisms in impulsivity and rodent models of addiction-like behaviours and thus the feasibility of treating impulsivity to remediate addiction.

Animal models of addiction-like behaviour

While many rodent paradigms are available to model addiction-related processes, including reinforcement mechanisms (Sanchis-Segura and Spanagel, 2006), we have limited our discussion to studies involving drug self-administration and reinstatement procedures in rodents. These behavioural paradigms typically involve operant responding for i.v., or in the case of alcohol, oral drug reward, often in the presence of contingent and non-contingent cues and/or contexts, and typically either fixed ratio (FR – where a fixed number of responses result in reward) or progressive ratio (PR – where an incremental number of responses result in reward) reinforcement schedules. Relapse-like behaviour can be assessed in a number of ways, either following extinction of the learned association between contingent cues and contexts in the absence of drug availability, or after a period of forced abstinence. Relapse is triggered by exposing subjects to cues or contexts previously associated with drug availability (cue/context induced), a physical or chemical stressor (e.g. foot-shock, yohimbine) or following the administration of a priming dose of drug (i.e. drug-primed). It should be noted that these addiction-related behaviours only model aspects of the addiction construct and fail to recapitulate for example, the patterns of compulsive drug-seeking and intake associated with addiction. It has been suggested that the generally poor translation of preclinical and laboratory findings to positive clinical outcomes may be related to the limitations of these models to more fully encompass real world human addiction (Haney and Spealman, 2008). While such animal behavioural models have been developed (e.g. Deroche-Gamonet *et al.*, 2004; Vanderschuren and Everitt, 2004; Belin *et al.*, 2008) only a few studies to date have assessed the ability of specific pharmacological agents to disrupt these behaviours (Pelloux *et al.*, 2012). Importantly, in terms of the clinical applicability of the current review, the subtypes of impulsivity discussed have been demonstrated to predict vulnerability to addiction as assessed by these contemporary rodent models (e.g. Belin *et al.*, 2008).

Pharmacological mechanisms of impulsivity: convergent mechanisms with addiction

Dopaminergic agents

Dysfunction of the midbrain dopaminergic systems has been implicated in several forms of impulsive behaviour (Pattij and Vanderschuren, 2008; Dalley and Roiser, 2012) and widely in the development and persistence of addiction (Volkow *et al.*, 2004; Thomas *et al.*, 2008; Sulzer, 2011; George *et al.*, 2012) not least as all drugs of abuse exert effects on the mesocorticolimbic dopamine neurons (Di Chiara and Imperato, 1988). However, the modulation of impulsivity and addiction-related behaviours by dopamine is complex and in the case of impulsivity often variable depending on the precise measure of impulsivity (Tables 1 and 2).

In keeping with the efficacy of stimulant medications in ADHD, acute administration of cocaine, amphetamine and

methylphenidate generally decreases impulsive choice in rats, increasing tolerance for delayed rewards in various discounting paradigms (Richards *et al.*, 1999; Wade *et al.*, 2000; Isles *et al.*, 2003; Winstanley *et al.*, 2003a; van Gaalen *et al.*, 2006a; Adriani *et al.*, 2007; Barbelivien *et al.*, 2008; Floresco *et al.*, 2008; Perry *et al.*, 2008; Krebs and Anderson, 2012). However, there have been notable exceptions (Evenden and Ryan, 1996; Cardinal *et al.*, 2000; Helms *et al.*, 2006; Stanis *et al.*, 2008a; Wooters and Bardo, 2011), which may reflect strain differences and paradigm-related effects. For example, methylphenidate reduces delay-discounting impulsivity in Wistar Kyoto rats, but has no effect in spontaneously hypertensive rats (Wooters and Bardo, 2011). Moreover, the presence of cues to signal delays can modulate the effects of stimulants in this task. Thus, amphetamine increases tolerance to cued delays (decreases impulsivity), but decreases tolerance to non-cued delays (Cardinal *et al.*, 2000). Stimulant medications also acutely increase impulsivity on tasks that assess action impulsivity; for example, the 5-CSRTT (Cole and Robbins, 1987; van Gaalen *et al.*, 2006b; Blondeau and Dellu-Hagedorn, 2007), but generally decrease impulsivity on the SSRTT (Feola *et al.*, 2000; Eagle and Robbins, 2003; Eagle *et al.*, 2007; 2009). These divergent effects lend support to the recently proposed dichotomy of 'waiting' versus 'stopping' forms of action impulsivity, which are differentially assayed by these two tasks (Dalley *et al.*, 2011).

Although stimulant drugs probably modulate impulsivity by enhancing dopaminergic tone, their effects are not always mirrored by drugs that act selectively through this mechanism, notably, dopamine re-uptake inhibitors (Table 1). Furthermore, the effect of amphetamine to reduce choice impulsivity is reduced in rats depleted of 5-HT (Winstanley *et al.*, 2003a). While these findings implicate additional neurotransmitters and mechanisms underlying the effects of indirect dopamine receptor agonists on impulsivity (including stimulants), the ability of directly acting D₁ and D₂-like receptor antagonists to oppose the effect of stimulants on impulsivity (e.g. van Gaalen *et al.*, 2006a,b) support a dopaminergic mechanism of action. Work over several years has localized these effects to specific corticostriatal sites (Table 2) with receptor subtype and region-specific effects. For example while oppositional effects of D₂-like receptor antagonism is observed between the nucleus accumbens (NAc) core and shell, decreasing and increasing impulsivity respectively in the 5-CSRTT (Besson *et al.*, 2010), a similar subregion-dependent distinction has not been reported for D₁ receptors (Pattij *et al.*, 2007a). It is unclear whether a similar divergence in dopaminergic modulation exists between the core and shell of the NAc with respect to impulsive choice; however, the effects of selective lesions of these subregions would appear to support this assertion (Ghods-Sharifi and Floresco, 2010). Further region-specific differentiation between D₁- and D₂-like receptor subtypes has been observed within the orbitofrontal cortex (OFC), with administration of a D₂/D₃, but not a D₁ receptor agonist found to reduce premature responding in highly impulsive (HI) animals (Winstanley *et al.*, 2010b). Similarly, administration of the D₂/D₃ receptor antagonist raclopride in the OFC increased impulsive choice, while a D₁ receptor antagonist had no effect (Pardey *et al.*, 2013). The significance of opponent interactions between NAc subregion and dopamine receptor subtype is unclear;

Table 1

Selected examples of the effects of acute systemic administration of dopaminergic compounds on measures of impulsivity and addiction-like behaviour in rodent models

Agent	Impulsive action		Impulsive choice		
	5-CSRTT	SSRTT	DD	Self-administration	Reinstatement
Dopamine re-uptake inhibitor/releaser					
Amphetamine	↑ (Harrison <i>et al.</i> , 1997; van Gaalen <i>et al.</i> , 2006b)	↓ (Eagle <i>et al.</i> , 2009)	↑ (Evenden and Ryan, 1996; Cardinal <i>et al.</i> , 2000) ↓ (Wade <i>et al.</i> , 2000; van Gaalen <i>et al.</i> , 2006a)	↑ ethanol (Pfeffer and Samson, 1985) ↑ FR cocaine (Barrett <i>et al.</i> , 2004; Xi <i>et al.</i> , 2009) ↓ PR for cocaine (Xi <i>et al.</i> , 2009)	↑ cocaine (De Vries <i>et al.</i> , 1998a)
Cocaine	↑ (van Gaalen <i>et al.</i> , 2006b)				↑ cocaine (De Vries <i>et al.</i> , 1998a)
Methylphenidate	↑ (Navarra <i>et al.</i> , 2008b; Milstein <i>et al.</i> , 2010; Pattij <i>et al.</i> , 2012) =§ (Fernando <i>et al.</i> , 2012)	↓§ (Eagle <i>et al.</i> , 2007)	↓ (van Gaalen <i>et al.</i> , 2006a)	↑ FR cocaine (Hiranita <i>et al.</i> , 2011) ↑ FR nicotine (Wooters <i>et al.</i> , 2008)	↑ drug-primed cocaine (Schenk and Partridge, 1999; Broos <i>et al.</i> , 2012a) = cue-induced cocaine (Economidou <i>et al.</i> , 2011)
GBR12909	↑ (van Gaalen <i>et al.</i> , 2006b)	= (Bari <i>et al.</i> , 2009)	↓ (van Gaalen <i>et al.</i> , 2006a)	↓ FR cocaine (Tella, 1995; Schenk, 2002) ↑ FR cocaine (Barrett <i>et al.</i> , 2004)	↑ drug-primed cocaine (Schenk, 2002)
D ₁ agonist					
SKF-81297	= (Winstanley <i>et al.</i> , 2010b)		= (Koffarnus <i>et al.</i> , 2011)	↓ FR ethanol (Cohen <i>et al.</i> , 1999) ↑ PR cocaine (Rowlett <i>et al.</i> , 2007)	↓ drug-primed cocaine (Self <i>et al.</i> , 1996)
SKF-82958				↓ FR cocaine (Caine <i>et al.</i> , 1999)	
D ₁ antagonist					
SCH-23390	↓ (Harrison <i>et al.</i> , 1997; Koskinen and Sirvio, 2001; van Gaalen <i>et al.</i> , 2006a,b)	= (Bari and Robbins, 2013)	↑* (amphetamine) (van Gaalen <i>et al.</i> , 2006a), = (Wade <i>et al.</i> , 2000)	↓ PR cocaine (Hubner and Moreton, 1991; Depoortere <i>et al.</i> , 1993) ↑ FR cocaine (Koob <i>et al.</i> , 1987)	↓ drug-primed cocaine (Self <i>et al.</i> , 1996; Schenk and Gittings, 2003) ↓ context-induced cocaine (Broos <i>et al.</i> , 2012a) ↓ drug-primed MA (Carati and Schenk, 2011) ↓ cue-induced nicotine (Liu <i>et al.</i> , 2010)

Table 1

Continued

Agent	Impulsive action		Impulsive choice		Reinstatement
	5-CSRTT	SSRTT	DD	Self-administration	
D ₂ agonist Quinpirole	↓§ (Fernando <i>et al.</i> , 2012)		= (Koffarnus <i>et al.</i> , 2011)	↓ FR cocaine (Caine and Koob, 1993) ↓ FR ethanol (Cohen <i>et al.</i> , 1998) = PR MA (Izzo <i>et al.</i> , 2001)	↑ drug-primed cocaine (Self <i>et al.</i> , 1996), ↑ cocaine (De Vries <i>et al.</i> , 2002)
D ₂ antagonist Eticlopride	= (van Gaalen <i>et al.</i> , 2006b) ↓* (amphetamine, cocaine, nicotine) (van Gaalen <i>et al.</i> , 2006b)		= (van Gaalen <i>et al.</i> , 2006a) ↑* (amphetamine) (van Gaalen <i>et al.</i> , 2006a)	↑ FR cocaine (Caine and Koob, 1994) ↓ PR cocaine (Ward <i>et al.</i> , 1996) ↓ PR MA (Izzo <i>et al.</i> , 2001)	↓ drug-primed cocaine (Schenk and Gittings, 2003) = drug-primed MA (Carati and Schenk, 2011) ↓ cue-induced nicotine (Liu <i>et al.</i> , 2010)
L741626	= (van Gaalen <i>et al.</i> , 2009) ↓* (amphetamine) (van Gaalen <i>et al.</i> , 2009)		= (Koffarnus <i>et al.</i> , 2011)		
Aripiprazole	↓§ (Besson <i>et al.</i> , 2010)			↑ FR cocaine (Roman <i>et al.</i> , 2013) ↓ PR MA (Wee <i>et al.</i> , 2007)	↓ cue-induced, drug-primed cocaine (Feltenstein <i>et al.</i> , 2007)
Sulpiride		= (Bari and Robbins, 2013)			
Haloperidol			= (Evenden and Ryan, 1996)		
Raclopride			↑ (Wade <i>et al.</i> , 2000)	↑ FR cocaine (Weissenborn <i>et al.</i> , 1996)	↓ drug-primed cocaine (Weissenborn <i>et al.</i> , 1996) ↓ context induced cocaine (Crombag <i>et al.</i> , 2002)
D ₁ /D ₂ antagonist Flupenthixol	↓* (DOI) (Koskinen and Sirvio, 2001)		↑ (Cardinal <i>et al.</i> , 2000; Wade <i>et al.</i> , 2000)	↑ FR, ↓ PR cocaine (Richardson <i>et al.</i> , 1994)	
D ₃ agonist 7-OH-PIPAT		= (Bari and Robbins, 2013)		↓ FR, ↑ PR cocaine (Caine and Koob, 1995)	= drug-primed cocaine (Khroyan <i>et al.</i> , 2000)
D ₃ antagonist Nafadotride	=§ (Besson <i>et al.</i> , 2010)	= (Bari and Robbins, 2013)			↓ cue-induced cocaine (Weiss <i>et al.</i> , 2001)

Table 1

Continued

Agent	Impulsive action		Impulsive choice		
	5-CSRTT	SSRTT	DD	Self-administration	Reinstatement
SB277011	= (van Gaalen <i>et al.</i> , 2009) = * (amphetamine) (van Gaalen <i>et al.</i> , 2009)			↓ FR, PR cocaine (Song <i>et al.</i> , 2012) ↓ PR MA (Higley <i>et al.</i> , 2011) ↓ PR nicotine (Ross <i>et al.</i> , 2007) = FR nicotine (Kameda <i>et al.</i> , 2000) = FR ethanol (Heidbreder <i>et al.</i> , 2007)	↓ drug-primed MA (Higley <i>et al.</i> , 2011) ↓ drug-primed ethanol (Heidbreder <i>et al.</i> , 2007)
Pramipexole			= (Koffarnus <i>et al.</i> , 2011)		
PG01037			= (Koffarnus <i>et al.</i> , 2011)		
D ₄ agonist					
PD168077		= (Bari and Robbins, 2013)			
D ₄ antagonist					
L745870	↓* (methylphenidate) (Milstein <i>et al.</i> , 2010)	= (Bari and Robbins, 2013)	= (Koffarnus <i>et al.</i> , 2011)		
ABT724			↑ (Koffarnus <i>et al.</i> , 2011)		

*Denotes significant effect on pharmacologically increased (5-CSRTT)/decreased levels of impulsivity, agent in parentheses.

§Denotes effect in selected high-impulsive rats.

↑, increased; ↓, decreased; =, no effect; DD, delay discounting; DOI, (+/-)-1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropan hydrochloride; MA, methamphetamine.

however, similar regional and neurochemical differentiation has been reported in the NAc of naturally occurring HI animals (Jupp *et al.*, 2013b; Simon *et al.*, 2013).

In keeping with an implicit role of enhanced dopaminergic neurotransmission in mediating drug reward and reinforcement, pharmacological agents affecting the dopaminergic systems have a range of effects on addiction-like behaviours in rodents. Systemic administration of D₁- and D₂-like receptor antagonists result in compensatory increases in responding for stimulant drugs under a FR schedule of reinforcement (e.g. Corrigall and Coen, 1991a; Barrett *et al.*, 2004), but decrease break points and thus motivation to work for drug under a PR schedule (e.g. Hubner and Moreton, 1991; Richardson *et al.*, 1994; Izzo *et al.*, 2001). By contrast, dopamine receptor agonists reduce responding for stimulant drugs under FR schedules, with D₂-like receptor agonists decreasing the rate of drug self-administration and D₁ receptor agonists increasing the latency to initiate this behaviour (Caine *et al.*, 1999); this distinction may reflect distinct roles of D₂ and D₁ receptors in drug reward and satiety (Suto and Wise, 2011). However, the effects of dopaminergic agents on opiate,

alcohol and nicotine self-administration are less clear with conflicting effects of pharmacological blockade in some cases (e.g. Dyr *et al.*, 1993; David *et al.*, 2006; Rowlett *et al.*, 2007). Nevertheless, dopamine receptor antagonists, given either systemically or directly in the NAc generally inhibit reinstatement of drug-seeking for all major drug classes (reviewed in Self, 2010). However, systemic D₁ receptor agonists fail to induce drug-seeking (Self *et al.*, 1996) and actually reduce both drug- and cue-induced relapse when given systemically (De Vries *et al.*, 1999). Thus, enhanced mesolimbic dopamine appears to trigger relapse through D₂ receptor mechanisms with D₂-like receptor agonists potentiating drug-seeking behaviour (Self *et al.*, 1996; Edwards *et al.*, 2007). Importantly, however, the effects of D₁ and D₂ receptor agonists, such as for impulsivity, appear again to be region-dependent, with D₁ receptor agonists in the NAc core and shell sufficient to stimulate drug relapse (Bachtell *et al.*, 2005), whereas D₂ receptor agonists increase relapse only when given in the NAc shell (Schmidt *et al.*, 2006). Similarly, while relapse/reinstatement can be blocked by the administration of D₁ receptor antagonists in the NAc shell (Bachtell *et al.*, 2005) and core

Table 2

Selected studies of region-specific dopaminergic interventions in impulsivity and addiction-related behaviour in rodent models

Agent	Region	Impulsive action		Impulsive choice		
		5-CSRTT	SSRTT	DD	Self-administration	Reinstatement
Dopamine re-uptake inhibitor/releaser						
Amphetamine	NAc	↑ (Cole and Robbins, 1987)			↑ FR ethanol (Samson <i>et al.</i> , 1993; 1999)	
Methylphenidate	NAc core	↑ (Economidou <i>et al.</i> , 2012)				
	NAc shell	= (Economidou <i>et al.</i> , 2012)				
D ₁ agonist						
SKF-38393	NAc	↑ (Pezze <i>et al.</i> , 2007)				
	NAc core					↑ cocaine (Bachtell <i>et al.</i> , 2005)
	NAc shell				= FR cocaine (Bachtell <i>et al.</i> , 2005)	↑ cocaine (Bachtell <i>et al.</i> , 2005)
	OFC	=§ (Winstanley <i>et al.</i> , 2010b)				
D ₁ antagonist						
SCH-23390	NAc core	↓ (Pattij <i>et al.</i> , 2007a)	= (Eagle <i>et al.</i> , 2011)		↓ PR cocaine (Bari and Pierce, 2005)	↓ context-induced ethanol (Chaudhri <i>et al.</i> , 2009)
	NAc shell	↓ (Pattij <i>et al.</i> , 2007a)			↓ PR cocaine (Bari and Pierce, 2005) ↑ FR cocaine (Maldonado <i>et al.</i> , 1993)	↓ drug-primed cocaine (Bachtell <i>et al.</i> , 2005)
	DS	↓ (Agnoli and Carli, 2011)	↓ (Eagle <i>et al.</i> , 2011)		↑ FR cocaine (Caine and Koob, 1995; Caine <i>et al.</i> , 1995)	
	PFC			↑ (Loos <i>et al.</i> , 2010; Pardey <i>et al.</i> , 2013)	↓ PR cocaine (Olsen and Duvauchelle, 2006)	↓ stress-induced cocaine (Capriles <i>et al.</i> , 2003) = drug-primed cocaine (Capriles <i>et al.</i> , 2003) ↓ drug-primed cocaine (Sun and Rebec, 2005)
	OFC	↓§ (Winstanley <i>et al.</i> , 2010b)		= (Pardey <i>et al.</i> , 2013)		↓ context-induced cocaine (Lasseter <i>et al.</i> , 2014) ↓ stress-induced cocaine (Capriles <i>et al.</i> , 2003)
	vHC			= (Abela and Chudasama, 2014)		

Table 2

Continued

Agent	Region	Impulsive action		Impulsive choice		
		5-CSRTT	SSRTT	DD	Self-administration	Reinstatement
D ₂ agonist						
7-OH-DPAT	NAc				= FR cocaine (Bachtell <i>et al.</i> , 2005)	
Quinpirole		↑ (Pezze <i>et al.</i> , 2007)			↑ FR ethanol (Samson and Chappell, 2003)	↑ cocaine (shell) (Schmidt <i>et al.</i> , 2006)
	OFC	↓§ (Winstanley <i>et al.</i> , 2010b)				
D ₂ antagonist						
Raclopride	OFC			↑ (Pardey <i>et al.</i> , 2013)		= stress, context-induced cocaine (Capriles <i>et al.</i> , 2003)
Sulpiride	DS		↑ (Eagle <i>et al.</i> , 2011)			
Eticlopride	NAc core	=* (amphetamine) (Pattij <i>et al.</i> , 2007a)			= FR cocaine (Bachtell <i>et al.</i> , 2005)	
Aripiprazole		=§ (Besson <i>et al.</i> , 2010)				
Sulpiride			= (Eagle <i>et al.</i> , 2011)			= drug-primed cocaine (Anderson <i>et al.</i> , 2006)
	NAc	= (Pezze <i>et al.</i> , 2007)			↑ FR ethanol (Levy <i>et al.</i> , 1991) ↑ FR cocaine, (Phillips <i>et al.</i> , 1994)	
	NAc shell					↓ drug-primed cocaine (Anderson <i>et al.</i> , 2006)
Aripiprazole		= § (Besson <i>et al.</i> , 2010)				
Eticlopride		=* (amphetamine) (Pattij <i>et al.</i> , 2007a)			↓ PR cocaine (Bari and Pierce, 2005) ↑ FR cocaine (Bachtell <i>et al.</i> , 2005)	↓ drug-primed cocaine (Bachtell <i>et al.</i> , 2005)
D ₃ antagonist						
Nafadotride	NAc core	↓§ (Besson <i>et al.</i> , 2010)				
	NAc shell	↑§ (Besson <i>et al.</i> , 2010)				
SB-277011A	NAc					↓ stress-induced cocaine (Xi <i>et al.</i> , 2004)

*Denotes effect on pharmacologically increased/decreased levels of impulsivity, agent in parentheses.

§Denotes an effect in selected high-impulsive rats.

↑, increased; ↓, decreased; =, no effect; DS, dorsal striatum; MA, methamphetamine; vHC, ventral hippocampus.

(Chaudhri *et al.*, 2009) the effects of D₂ receptor antagonists are limited to the NAc shell (Anderson *et al.*, 2006).

The findings reviewed earlier, especially in relation to psychostimulants, are consistent with the effects of acute systemically administered dopaminergic agents on impulsive action (e.g. as measured in the 5-CSRTT), but not impulsive choice. Specifically, dopamine compounds that reduce or increase impulsive action also, in general, reduce or increase measures of drug reward and relapse (Table 1), with convergence in terms of both regional and dopamine receptor subtype specificity (Table 2). However, although systemically administered D₁ and D₂ receptor agonists enhance reward and at least in the case of D₂ receptors, stimulate relapse, dopamine receptor agonists either have no effect (D₁-like; Winstanley *et al.*, 2010b) or decrease premature responding in the 5-CSRTT (D₂-like; Besson *et al.*, 2010). Moreover, D₂-like receptor blockade in the NAc shell reduces drug reward and reinstatement (Anderson *et al.*, 2006), but increases premature responding in the 5-CSRTT (Besson *et al.*, 2010). Thus, manipulations that acutely alter dopamine receptor function appear unlikely to collectively suppress impulsivity and addiction-like behaviours.

5-Hydroxytryptaminergic agents

There is widespread support for a significant role of 5-HT in modulating impulsivity and addiction-related processes (Kranz *et al.*, 2010; Hayes and Greenshaw, 2011; Kirby *et al.*, 2011; Dalley and Roiser, 2012; Miyazaki *et al.*, 2012), through interactions with dopamine (Kapur and Remington, 1996; Di Matteo *et al.*, 2008) and other neurotransmitter systems (Fink and Gothert, 2007). A summary of the involvement of different 5-HT receptor subtypes in impulsivity and their putative loci of action is given in Tables 3 and 4. Near-complete depletion of 5-HT in the brain increases impulsivity in the 5-CSRTT (Harrison *et al.*, 1997; Winstanley *et al.*, 2004), the Go/No-go task (Harrison *et al.*, 1999) and the SSRTT (Eagle *et al.*, 2009). However, 5-HT depletion has variable effects on delay-discounting impulsivity, which may be due to differing experimental protocols (for discussion, see Winstanley *et al.*, 2006). Similarly, enhancing 5-HT levels via systemically administered 5-HT re-uptake inhibitors reduces impulsivity in the 5-CSRTT (Baarendse and Vanderschuren, 2012; Humpston *et al.*, 2013), but has no effect on SSRTT performance (Bari *et al.*, 2009) or delay discounting (Evenden and Ryan, 1996; Baarendse and Vanderschuren, 2012). Studies of receptor selective ligands have predominantly implicated 5-HT₂ receptors in mediating these effects. 5-HT_{2C} receptor antagonists such as SB242084 increase premature responding in the 5-CSRTT (Winstanley *et al.*, 2004; Fletcher *et al.*, 2007) while the 5-HT_{2C} receptor agonist WAY163909 reduces impulsivity on this task (Navarra *et al.*, 2008a). Interestingly, the mixed 5-HT_{2A/2C} receptor antagonist ketanserin reduced (Passetti *et al.*, 2003; Talpos *et al.*, 2006; Fletcher *et al.*, 2007), while the 5-HT_{2A/2C} receptor agonist DOI ((+/-)-1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropan hydrochloride) increased premature responding in the 5-CSRTT (Koskinen *et al.*, 2000). Consistent with these results the selective 5-HT_{2A} receptor antagonist M100907 decreased premature responding (Winstanley *et al.*, 2003b; Fletcher *et al.*, 2007), similar to the effect of the mixed 5-HT_{2B/C} receptor agonist Ro60-175 (Fletcher *et al.*, 2007). It is unclear whether delay discounting is also bi-directionally modulated

by 5-HT_{2A} and 5-HT_{2C} receptors, but both ketanserin and the 5-HT_{2C} receptor antagonist SB242084 decrease this measure of impulsivity (Hadamitzky and Koch, 2009; Hadamitzky *et al.*, 2009; Paterson *et al.*, 2012). Finally, 5-HT₁ receptors have been implicated in several variants of impulsivity, but the results have been quite mixed. Thus, activation of 5-HT_{1A} receptors increases impulsivity on the delay-discounting task (Winstanley *et al.*, 2005b; van den Bergh *et al.*, 2006; Stanis *et al.*, 2008b; Blasio *et al.*, 2012) and either increases (Carli and Samanin, 2000), decreases (Blokland *et al.*, 2005) or has no effect (Winstanley *et al.*, 2003b; Carli *et al.*, 2006) on premature responding in the 5-CSRTT.

Wide-ranging evidence also implicates a role for 5-HT in reward and addiction (reviewed Filip *et al.*, 2010; Hayes and Greenshaw, 2011; Kirby *et al.*, 2011). However, owing to the complexity of this neurotransmitter system and its influence on dopamine and other neurotransmitter systems, the precise role of 5-HT in addiction-related processes is not always clear. Globally depleting 5-HT in the brain tends to potentiate drug reward, increasing stimulant drug self-administration under both FR and PR schedules (e.g. Lyness *et al.*, 1980; Roberts *et al.*, 1994) and drug-primed relapse (Tran-Nguyen *et al.*, 2001). Conversely, selective 5-HT re-uptake inhibitors generally decrease drug self-administration and relapse (Glatz *et al.*, 2002; Simon O'Brien *et al.*, 2011).

A survey of specific 5-HT receptor subtypes reveals that 5-HT_{1A} receptor activation enhances the rewarding effects of drugs by decreasing drug self-administration under FR schedules while increasing PR responding for cocaine (Peltier and Schenk, 1993; Parsons *et al.*, 1998) and alcohol (Wilson *et al.*, 1996) reinforcement. Tonic activity at 5-HT_{1A} receptors appears important for the rewarding effects of stimulant drugs, demonstrated by the inhibition of drug-primed reinstatement of cocaine-seeking by the selective 5-HT_{1A} receptor antagonist WAY100635 (Schenk, 2000; Burmeister *et al.*, 2004). Conversely, activation of 5-HT_{1B} receptors attenuates drug reward, increasing FR and decreasing PR responding for cocaine (Parsons *et al.*, 1998). Again, tonic activity at this receptor subtype appears important for cocaine-motivated behaviours, including reinstatement (David *et al.*, 2004; Przegalinski *et al.*, 2008).

However, the role of 5-HT₂ receptors in addiction-like behaviour is more nuanced. Thus, while systemically administered 5-HT_{2A} receptor antagonists appear to have no effect on the self-administration of cocaine or nicotine (Fletcher *et al.*, 2002; 2012; Nic Dhonnchadha *et al.*, 2009) they do reduce responding for alcohol when infused into the ventral tegmental area (Ding *et al.*, 2009). Selective 5-HT_{2A} receptor antagonists also inhibit drug- and cue-primed nicotine- and cocaine-seeking (Nic Dhonnchadha *et al.*, 2009; Fletcher *et al.*, 2012). The differential effect of these compounds on self-administration and relapse has been suggested to relate to the inhibition of dopamine release in the dorsal striatum, but not the NAc by these agents following drug administration (Murnane *et al.*, 2013). By contrast, selective 5-HT_{2C} receptor antagonists tend to increase rates of cocaine (Fletcher *et al.*, 2002) and alcohol (Tomkins *et al.*, 2002) self-administration, and drug-primed relapse (Burmeister *et al.*, 2004). Consistent with these findings, selective 5-HT_{2C} receptor agonists reduce self-administration and cue-induced cocaine (Cunningham *et al.*, 2011) and nicotine (Higgins *et al.*, 2012) reinstatement,

Table 3

Selected studies employing 5-hydroxytryptaminergic interventions in models of impulsivity and addiction-related behaviour in rodent models

Agent	Impulsive action		Impulsive choice		Reinstatement
	5-CSRTT	SSRTT	DD	Self-administration	
5-HT re-uptake inhibitor/releaser					
Citalopram	↓ (Baarendse and Vanderschuren, 2012; Humpston <i>et al.</i> , 2013)	= (Bari <i>et al.</i> , 2009)	= (Evenden and Ryan, 1996; Baarendse and Vanderschuren, 2012)	= FR cocaine (Hiranita <i>et al.</i> , 2009)	↓ drug-primed cocaine (Ruedi-Bettschen <i>et al.</i> , 2010)
Fluvoxamine	= (Tsutsui-Kimura <i>et al.</i> , 2009)			↓ FR ethanol (Lamb and Jarbe, 2001)	
Fluoxetine	↓ (Humpston <i>et al.</i> , 2013)			↓ FR ethanol (Simon O'Brien <i>et al.</i> , 2011) ↓ FR cocaine (Glatz <i>et al.</i> , 2002)	↓ stress-induced ethanol (Simon O'Brien <i>et al.</i> , 2011)
Paroxetine	↓ (Humpston <i>et al.</i> , 2013)				
5-HT depletion					
5,7-dihydroxytryptamine	↑ (Harrison <i>et al.</i> , 1997; Winstanley <i>et al.</i> , 2004)		↑ (Wogar <i>et al.</i> , 1993; Bizot <i>et al.</i> , 1999; Mobini <i>et al.</i> , 2000) = (Winstanley <i>et al.</i> , 2003a; 2004)	↑ FR MA (Fletcher <i>et al.</i> , 1999) ↑ PR cocaine (Roberts <i>et al.</i> , 1994)	↓ cue-induced cocaine (Tran-Nguyen <i>et al.</i> , 2001) ↑ drug-primed cocaine (Tran-Nguyen <i>et al.</i> , 2001)
5-HT _{1A} agonist					
8-OH-DPAT	↑ (Carli and Samanin, 2000) = (Winstanley <i>et al.</i> , 2003b) ↓ (Blokland <i>et al.</i> , 2005)		↑ (Winstanley <i>et al.</i> , 2005b; Stanis <i>et al.</i> , 2008a; Blasio <i>et al.</i> , 2012)	↓ FR cocaine (Peltier and Schenk, 1993) ↓ PR cocaine (Parsons <i>et al.</i> , 1998) ↓ FR ethanol (Wilson <i>et al.</i> , 1996)	
Flesinoxan			↑ (van den Bergh <i>et al.</i> , 2006)		
5-HT _{1A} antagonist					
WAY100635	=/≠ * (methylphenidate) (Milstein <i>et al.</i> , 2010)				↓ drug-primed cocaine (Schenk, 2000; Burmeister <i>et al.</i> , 2004)
m-MPPI				↓ PR cocaine (Parsons <i>et al.</i> , 1998)	
5-HT _{1B} antagonist					
GR55562	=/≠ * (methylphenidate) (Milstein <i>et al.</i> , 2010)				
GR127935			= (van den Bergh <i>et al.</i> , 2006)		↓ drug-primed cocaine (Przegalinski <i>et al.</i> , 2008), ↓ cue-induced cocaine (Przegalinski <i>et al.</i> , 2008)

Table 3

Continued

Agent	Impulsive action		Impulsive choice		
	5-CSRTT	SSRTT	DD	Self-administration	Reinstatement
5-HT _{1A/1B} agonist					
Etoprazine			↓ (van den Bergh <i>et al.</i> , 2006)		
RU24969	= (Evenden, 1999b)			↓ FR, ↑ PR cocaine (Parsons <i>et al.</i> , 1998)	↓ drug-primed cocaine (Acosta <i>et al.</i> , 2005) ↓ cue-induced cocaine (Acosta <i>et al.</i> , 2005)
5-HT _{2A} antagonist					
M100907	↓ (Winstanley <i>et al.</i> , 2003b; Fletcher <i>et al.</i> , 2007) ↓* (dizocilpine) (Higgins <i>et al.</i> , 2003) ↓* (5,7-dihydroxytryptamine) (Winstanley <i>et al.</i> , 2004)			= FR cocaine (Nic Dhonnchadha <i>et al.</i> , 2009) = PR cocaine (Fletcher <i>et al.</i> , 2002) = FR, PR nicotine (Fletcher <i>et al.</i> , 2012)	↓ cue-induced cocaine (Nic Dhonnchadha <i>et al.</i> , 2009) ↓ drug- primed nicotine, cue-induced nicotine (Fletcher <i>et al.</i> , 2012)
5-HT _{2A/C} agonist					
DOI	↑ (Koskinen <i>et al.</i> , 2000; Blokland <i>et al.</i> , 2005)		↑ (Hadamitzky and Koch, 2009; Hadamitzky <i>et al.</i> , 2009; Blasio <i>et al.</i> , 2012)	↓ FR ethanol (Maurel <i>et al.</i> , 1999)	
5-HT _{2A/C} antagonist					
Ketanserin	↓ (Passetti <i>et al.</i> , 2003; Talpos <i>et al.</i> , 2006; Fletcher <i>et al.</i> , 2007) ↓* (DOI) (Koskinen <i>et al.</i> , 2000)		↓* (DOI) (Hadamitzky and Koch, 2009; Hadamitzky <i>et al.</i> , 2009) = (Paterson <i>et al.</i> , 2012 22094071)	↓ FR nicotine (Levin <i>et al.</i> , 2008)	↓ cue-induced cocaine (Burmeister <i>et al.</i> , 2004)
Ritanserin	↓* (DOI) (Koskinen <i>et al.</i> , 2000)			= FR MA (Fletcher, 1998)	= cocaine (Schenk, 2000)
5-HT _{2C} agonist					
WAY163909	↓ (Navarra <i>et al.</i> , 2008a)			↓ FR cocaine (Cunningham <i>et al.</i> , 2011)	↓ cue-induced cocaine (Cunningham <i>et al.</i> , 2011)
Lorcaserin				↓ FR nicotine (Higgins <i>et al.</i> , 2012)	↓ nicotine, cue-induced nicotine (Higgins <i>et al.</i> , 2012)
5-HT _{2C} antagonist					
SB242084	↑ (Winstanley <i>et al.</i> , 2004; Fletcher <i>et al.</i> , 2007)		↓ (Paterson <i>et al.</i> , 2012)	↑ FR, PR cocaine (Fletcher <i>et al.</i> , 2002) ↑ FR ethanol (Tomkins <i>et al.</i> , 2002)	= cue-induced MA (Graves and Napier, 2012) ↑ drug-primed cocaine, (Burmeister <i>et al.</i> , 2004)

Table 3

Continued

Agent	Impulsive action		Impulsive choice		Reinstatement
	5-CSRTT	SSRTT	DD	Self-administration	
5-HT _{2B/C} agonist Ro60-0175	↓ (Fletcher <i>et al.</i> , 2007)			↓ FR, PR cocaine (Fletcher <i>et al.</i> , 2004) ↓ FR, PR nicotine (Fletcher <i>et al.</i> , 2012)	↓ cue-induced nicotine (Fletcher <i>et al.</i> , 2012) ↓ stress- induced cocaine (Fletcher <i>et al.</i> , 2008) ↓ context- induced cocaine (Fletcher <i>et al.</i> , 2008)
5-HT _{2B/C} antagonist SER-082	= (Talpos <i>et al.</i> , 2006)		↓ (Talpos <i>et al.</i> , 2006)	= FR cocaine (Filip, 2005)	= cue-induced MA (Graves and Napier, 2012) = drug-primed cocaine (Filip, 2005) = cue-induced cocaine (Filip, 2005)
5-HT _{2B} antagonist SB215505	= (Fletcher <i>et al.</i> , 2007)			↓ FR, ↑PR cocaine (Fletcher <i>et al.</i> , 2002)	↑ cue-induced cocaine (Fletcher <i>et al.</i> , 2002)
5-HT ₃ antagonist MDL72222	↓ (Evenden, 1999a)			↓ FR ethanol (McKinzie <i>et al.</i> , 2000) = FR nicotine (Corrigall and Coen, 1994) = PR cocaine (Lacosta and Roberts, 1993)	
Ondansetron				= FR cocaine (Lane <i>et al.</i> , 1992) ↓ PR cocaine (Davidson <i>et al.</i> , 2002)	↓ stress- induced ethanol (Le <i>et al.</i> , 2006)
5-HT ₆ antagonist SB270146A	= (Talpos <i>et al.</i> , 2006)		= (Talpos <i>et al.</i> , 2006)		
CMP42	= (de Bruin <i>et al.</i> , 2013)			↓ FR nicotine (de Bruin <i>et al.</i> , 2013)	↓ cue- induced nicotine (de Bruin <i>et al.</i> , 2013) ↓ cue- induced ethanol (de Bruin <i>et al.</i> , 2013)

*Denotes effect on pharmacologically increased/decreased levels of impulsivity, agent in parentheses.

↑, increased; ↓, decreased; =, no effect; DOI, (+/-)-1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropan hydrochloride; MA, methamphetamine.

an effect that appears to be mediated by the prefrontal cortex (PFC; Pentkowski *et al.*, 2010). Selective 5-HT_{2B} receptor antagonism enhances drug reward, reducing responding for cocaine under a FR schedule, while increasing responding under a PR schedule and augmenting drug-primed relapse behaviour (Fletcher *et al.*, 2002). However, mixed 5-HT_{2B/C}

receptor antagonists have no effect on drug self-administration or drug-seeking behaviour (Filip, 2005; Graves and Napier, 2012), although mixed 5-HT_{2B/C} receptor agonists decreased these behaviours (Fletcher *et al.*, 2004; 2008; 2012). Finally, 5-HT₃ receptor antagonism generally reduces responding for ethanol and cocaine (Tomkins *et al.*, 1995;

Table 4

Selected studies of region-specific 5-hydroxytryptaminergic interventions in rodent models of impulsivity

	Region	Impulsive action	Impulsive choice
Intervention		5-CSRTT	DD
5-HT depletion			
5,7-dihydroxytryptamine	PFC	= (Fletcher <i>et al.</i> , 2009)	
	NAC	= (Fletcher <i>et al.</i> , 2009)	
5-HT _{1A} agonist			
8-OH-DPAT	PFC	= (Winstanley <i>et al.</i> , 2003b) =* (m-CPP) (Carli <i>et al.</i> , 2006)	
5-HT _{2A} antagonist			
M100907	PFC	↓ (Winstanley <i>et al.</i> , 2003b) = (Robinson <i>et al.</i> , 2008a) ↓* (m-CPP) (Carli <i>et al.</i> , 2006)	
	DS	↓* (m-CPP) (Agnoli and Carli, 2012)	
	NAC	↓ (Robinson <i>et al.</i> , 2008a)	
5-HT _{2C} antagonist			
SB242084	NAC	↑ (Robinson <i>et al.</i> , 2008a)	
	PFC	= (Robinson <i>et al.</i> , 2008a)	
5-HT _{2A/C} agonist			
DOI	NAC Core	= (Koskinen and Sirvio, 2001)	
	NAC Shell	= (Koskinen and Sirvio, 2001)	
	OFC	= (Hadamitzky and Koch, 2009)	↑ (Wischhof <i>et al.</i> , 2011)
	BLA	= (Hadamitzky and Koch, 2009)	
	OFC + BLA	↑ (Hadamitzky and Koch, 2009)	
5-HT _{2A/C} antagonist			
Ketanserin	PFC	↓ (Passetti <i>et al.</i> , 2003)	
	OFC	= (Hadamitzky and Koch, 2009)	
	BLA	= (Hadamitzky and Koch, 2009)	
	OFC + BLA	= (Hadamitzky and Koch, 2009)	
5-HT _{2B/C} agonist			
Ro60-0175	DS	↓* (m-CPP) (Agnoli and Carli, 2012)	

*Denotes effect on pharmacologically increased/decreased levels of impulsivity, agent in parentheses.

↑, increased; ↓, decreased; =, no effect; BLA, basolateral amygdala; m-CPP, meta-chloropirazine; DOI, (+/-)-1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropan hydrochloride; DS, dorsal striatum.

McKinzie *et al.*, 2000; Davidson *et al.*, 2002), similar to the effects of 5-HT₆ receptor antagonists, which decrease nicotine self-administration as well as both cue- and drug-primed relapse (de Bruin *et al.*, 2013).

The findings reviewed earlier indicate a significant overlap in 5-hydroxytryptaminergic mechanisms affecting impulsivity and addiction-like behaviours in rodents (Table 3). Again, similar to the effect at other transmitter systems, pharmacological agents that reduce impulsive action (rather than impulsive choice) also reduce drug self-administration and drug-seeking, whereas agents that enhance impulsive action also potentiate addiction-like behaviours. However, despite an involvement of 5-HT_{1A}, 5-HT_{1B} and 5-HT₆ receptors in reward and addiction-like behaviour (reviewed earlier), there is no convincing evidence they play a major role in impulsive action.

Noradrenergic agents

A role for noradrenaline in impulsivity is substantiated by the clinical efficacy of amphetamine and methylphenidate in ADHD, which act to enhance noradrenaline as well as dopamine transmission in the brain [reviewed in Del Campo *et al.* (2011)] and more specifically by the effectiveness of the selective noradrenaline re-uptake inhibitor (NARI) atomoxetine in ADHD (Simpson and Plosker, 2004; Faraone *et al.*, 2005) and animal models of impulsivity (Blondeau and Dellu-Hagedorn, 2007; Navarra *et al.*, 2008b; Robinson *et al.*, 2008b; Tsutsui-Kimura *et al.*, 2009; Fernando *et al.*, 2012; see Table 5). In addition, mixed 5-HT/noradrenaline re-uptake inhibitors such as desipramine (van Gaalen *et al.*, 2006b; Paine *et al.*, 2007), milnacipran (Tsutsui-Kimura *et al.*, 2009) and venlafaxine (Humpston *et al.*, 2013) are effective in

Table 5

Selected studies of systemically administered (unless otherwise stated) noradrenergic interventions in rodent models of impulsivity and addiction-like behaviour

Agent	Impulsive action		Impulsive choice		
	5-CSRTT	SSRTT	DD	Self-administration	Reinstatement
NA re-uptake inhibitor					
Atomoxetine	↓§ (Blondeau and Dellu-Hagedorn, 2007; Fernando <i>et al.</i> , 2012) ↓ (Navarra <i>et al.</i> , 2008b; Robinson <i>et al.</i> , 2008b; Tsutsui-Kimura <i>et al.</i> , 2009)	↓ (Robinson <i>et al.</i> , 2008b; Bari <i>et al.</i> , 2009)	↓ (Robinson <i>et al.</i> , 2008b) = (Baarendse and Vanderschuren, 2012) ↑ (Broos <i>et al.</i> , 2012b)		↓ cue-induced cocaine (Economidou <i>et al.</i> , 2009; 2011)
Reboxetine	↓ (Liu <i>et al.</i> , 2009)			↓ FR nicotine (Rauhut <i>et al.</i> , 2002)	
NA/DA re-uptake inhibitor					
Bupropion	= (Humpston <i>et al.</i> , 2013)			↓ FR MA (Reichel <i>et al.</i> , 2009) ↓ FR nicotine (Liu <i>et al.</i> , 2008) = PR nicotine (Bruijnzeel and Markou, 2003)	↑ cue-induced nicotine (Liu <i>et al.</i> , 2008)
NA/5-HT re-uptake inhibitor					
Desipramine	↓ (van Gaalen <i>et al.</i> , 2006b; Paine <i>et al.</i> , 2007; Pattij <i>et al.</i> , 2012)		= (van Gaalen <i>et al.</i> , 2006a)	= FR cocaine (Tella, 1995)	↓ cue-induced ethanol (Simon O'Brien <i>et al.</i> , 2011)
Milnacipran	↓ (Tsutsui-Kimura <i>et al.</i> , 2009)				↓ cue-induced ethanol (Simon O'Brien <i>et al.</i> , 2011)
Subutramine	= (Humpston <i>et al.</i> , 2013)				
Venlafaxine	↓ (Humpston <i>et al.</i> , 2013)				
α ₁ agonist					
Phenylephrine			= (van Gaalen <i>et al.</i> , 2006a)		
α ₁ antagonist					
Prazosin	↓ (Liu <i>et al.</i> , 2009) ↓* (DOI) (Koskinen <i>et al.</i> , 2003)			= FR cocaine (Ecke <i>et al.</i> , 2012) ↓ FR nicotine (Forget <i>et al.</i> , 2010)	↓ stress- induced ethanol (Le <i>et al.</i> , 2011) ↓ drug- primed cocaine (Zhang and Kosten, 2005) ↓ drug-primed, cue-induced nicotine (Forget <i>et al.</i> , 2010)

Table 5

Continued

Agent	Impulsive action		Impulsive choice		Self-administration	Reinstatement
	5-CSRTT	SSRTT	DD			
α_2 agonist						
Guanfacine	↓§ (Fernando <i>et al.</i> , 2012) ↓ (Milstein <i>et al.</i> , 2007)		↓ vHC (Abela and Chudasama, 2014)			↓ stress-induced cocaine (Buffalari <i>et al.</i> , 2012) ↓ stress- induced ethanol (Le <i>et al.</i> , 2011) ↓ cue-induced cocaine (Smith and Aston-Jones, 2011)
Clonidine			↑ (van Gaalen <i>et al.</i> , 2006a)			↓ stress- induced cocaine (Erb <i>et al.</i> , 2000) = drug primed cocaine (Erb <i>et al.</i> , 2000) ↓ stress- induced nicotine (Zislis <i>et al.</i> , 2007)
Lofexidine					↓ FR ethanol (Le <i>et al.</i> , 2005)	↓ stress- induced ethanol (Le <i>et al.</i> , 2005)
Imidazoline						↓ cue- induced cocaine (Smith and Aston-Jones, 2011)
α_2 antagonist						
Yohimbine	↑ (Sun <i>et al.</i> , 2010; Torregrossa <i>et al.</i> , 2012)				↑ FR ethanol (Le <i>et al.</i> , 2005) ↑FR, PR nicotine (Li <i>et al.</i> , 2012)	↑ cocaine (Feltenstein and See, 2006) ↑ MA (Shepard <i>et al.</i> , 2004) ↑ ethanol (Le <i>et al.</i> , 2005) ↑ nicotine (Feltenstein <i>et al.</i> , 2012) ↑ cue- induced cocaine (Feltenstein <i>et al.</i> , 2011; Buffalari <i>et al.</i> , 2012)
Atipamezole	↑ (Koskinen <i>et al.</i> , 2003)					
Idazoxan	= (Humpston <i>et al.</i> , 2013)					
β_2 agonist						
Clenbuterol	↓ (Pattij <i>et al.</i> , 2012)					
β antagonist						
Propranolol	↓* (methylphenidate) (Milstein <i>et al.</i> , 2010)				↓ FR, PR ethanol (Gilpin and Koob, 2010) ↓ FR cocaine (Harris <i>et al.</i> , 1996)	↓ cue- induced nicotine (Chiamulera <i>et al.</i> , 2010)

*Denotes effect on pharmacologically increased/decreased levels of impulsivity, agent in parentheses.

§Denotes an effect in selected high-impulsive rats.

↑, increased; ↓ decreased; = no effect; MA, methamphetamine; vHC, ventral hippocampus.

reducing action impulsivity. These findings contrast with the mixed dopamine/noradrenaline re-uptake inhibitor bupropion, which has no effect on impulsive action (Humpston *et al.*, 2013) presumably as a result of opponent effects of dopamine on this task (Dalley *et al.*, 2011). The effects of noradrenaline on impulsive action appear to be mediated by α_1 -, α_2 - and β -adrenoceptors. Thus, systemic administration of α_1 receptor antagonists (Milstein *et al.*, 2007; Liu *et al.*, 2009) and α_2 -adrenoceptor agonists (Milstein *et al.*, 2007; Fernando *et al.*, 2012) reduce, whereas α_2 -adrenoceptor antagonists, increase action impulsivity (Koskinen *et al.*, 2003; Sun *et al.*, 2010; Torregrossa *et al.*, 2012). An action at β -adrenoceptors has also been implicated in this form of impulsivity, with a recent study demonstrating an effect of the β_2 -adrenoceptor selective agonist clenbuterol to reduce impulsivity, as measured by the 5-CSRTT (Pattij *et al.*, 2012). This appears somewhat at odds with data demonstrating an effect of the non-selective β -adrenoceptor antagonist propranolol to reduce premature responding induced by methylphenidate (Milstein *et al.*, 2010). Clearly further studies are required to delineate the relative contributions of β_1 - and β_2 -adrenoceptors to this behaviour.

While there is conflicting evidence for a role of NARIs in impulsive choice (van Gaalen *et al.*, 2006a; Robinson *et al.*, 2008b; Baarendse and Vanderschuren, 2012; Broos *et al.*, 2012b), systemic administration of the α_2 -adrenoceptor agonist clonidine appears to increase this form of impulsivity (van Gaalen *et al.*, 2006a), but curiously has the opposite effect when infused directly in the hippocampus (Abela and Chudasama, 2014). Consistent with region-specific effects of noradrenaline transmission on distinct impulsivity subtypes a recent study found that atomoxetine reduces impulsivity in the 5-CSRTT when administered in the NAc shell, but not the NAc core or the PFC (Economidou *et al.*, 2012).

Noradrenaline is thought to contribute to drug reward and addiction through dopamine-dependent and dopamine-independent processes (reviewed in Weinshenker and Schroeder, 2007). Pharmacological manipulation of this system affects several measures of addiction-related behaviours in rodents (Table 5). While these generally have little effect on psychostimulant self-administration (reviewed in Sofuoglu and Sewell, 2009) there is mounting evidence that noradrenergic mechanisms modulate ethanol, nicotine and opiate self-administration, specifically *via* α_1 - (Rasmussen *et al.*, 2009; Forget *et al.*, 2010), but not α_2 - (Le *et al.*, 2005; Marinelli *et al.*, 2007; Li *et al.*, 2014) adrenoceptors. Thus, α_1 -adrenoceptor antagonists such as prazosin attenuate drug-primed reinstatement (Zhang and Kosten, 2005), putatively by decreasing dopamine release in the NAc (Lane *et al.*, 1988).

Previous research indicates that noradrenaline plays a critical role in cue- and stress-induced drug relapse. Atomoxetine has been found to inhibit cue-induced cocaine reinstatement (Economidou *et al.*, 2009; 2011), while the mixed 5-HT/noradrenaline re-uptake inhibitors desipramine and milnacipran reduce cue-induced ethanol reinstatement (Simon O'Brien *et al.*, 2011). Conversely, the mixed noradrenaline/dopamine re-uptake inhibitor bupropion *enhances* nicotine cue-induced reinstatement (Liu *et al.*, 2008) presumably as a consequence of increased tonic activity at dopamine receptors. In terms of specific receptor subtypes α_1 -adrenoceptor antagonists (e.g. Forget *et al.*, 2010; Le *et al.*,

2011) and α_2 -adrenoceptor agonists (e.g. Erb *et al.*, 2000; Le *et al.*, 2005; Zislin *et al.*, 2007; Smith and Aston-Jones, 2011) reduce reinstatement to drug-associated cues and various stressors. By contrast, the α_2 -adrenoceptor antagonist yohimbine facilitates reinstatement to all major classes of abused drugs (psychostimulants, alcohol, opiates, nicotine) (Shepard *et al.*, 2004; Le *et al.*, 2005; Feltenstein and See, 2006; Banna *et al.*, 2010; Feltenstein *et al.*, 2012). Yohimbine also potentiates the effects of cue-exposure on cocaine-seeking (Feltenstein *et al.*, 2011). In addition, β -adrenoceptors contribute to cue- and stress-induced reinstatement (e.g. Leri *et al.*, 2002; Chiamulera *et al.*, 2010).

In keeping with the findings reviewed to date, there is considerable overlap in noradrenergic manipulations of impulsive action and addiction-like behaviour. Thus, drugs that decrease action impulsivity also act to reduce drug self-administration and/or inhibit relapse and *vice versa*. However, there is little overlap of these behaviours with impulsive choice (Table 5).

Glutamatergic agents

As shown in Table 6, systemic administration of non-competitive glutamatergic NMDA receptor antagonists predominantly have the effect of increasing impulsive action (Amitai *et al.*, 2007; Paine *et al.*, 2007; Oliver *et al.*, 2009; Fletcher *et al.*, 2011) and impulsive choice (Floresco *et al.*, 2008; Cottone *et al.*, 2013). At least in the case of impulsive action this effect appears to be mediated by GluN2B-containing receptors (for nomenclature see Alexander *et al.*, 2013b). Thus, the selective GluN2B NMDA receptor antagonist Ro63-1908 decreased impulsivity in the 5-CSRTT (Higgins *et al.*, 2005; Burton and Fletcher, 2012). Qualitatively similar effects are reported for metabotropic glutamate receptor (mGlu) antagonists (for nomenclature see Alexander *et al.*, 2013a), specifically acting at the mGlu₅ subtype on the 5-CSRTT (Semenova and Markou, 2007). Interestingly, mGlu₁ receptor antagonism decreased impulsive choice on a delay-discounting task (Sukhotina *et al.*, 2008).

While there is an abundance of evidence implicating NMDA, AMPA, kainic acid receptors, metabotropic receptors [e.g. mGlu₇ (Li *et al.*, 2010)] and excitatory amino acid transporters in addiction-related behaviours (for a comprehensive review, see Gass and Olive, 2008), we have limited our discussion to only those glutamatergic agents evaluated in rodent models of impulsivity. NMDA receptor antagonists generally attenuate drug self-administration and measures of drug reward including conditioned-place preference (e.g. Hyttia *et al.*, 1999; Glick *et al.*, 2001; Blokhina *et al.*, 2005; Yonghui *et al.*, 2006; Sabino *et al.*, 2013). One notable exception, however, is MK-801, which increases cocaine self-administration (Allen *et al.*, 2005) and augments relapse to cocaine (De Vries *et al.*, 1998b). Curiously, MK-801 also inhibits cue-induced relapse to ethanol (von der Goltz *et al.*, 2009), an effect resembling the actions of selective GluN2B NMDA receptor antagonists to reduce drug- and cue-induced reinstatement of ethanol (Vengeliene *et al.*, 2005; Wang *et al.*, 2010) and nicotine seeking (Gipson *et al.*, 2013). Systemically administered GluN2B receptor antagonists also reduce the self-administration of ethanol in rats (Wang *et al.*, 2010). Other studies have shown that mGlu₁, mGlu_{2/3} and mGlu₅ receptor antagonists reduce self-administration of, and

Table 6

Selected studies of systemically administered (unless otherwise stated) interventions of glutamatergic, GABAergic, opioidergic, cholinergic and cannabinoid neurotransmission in rodent models of impulsivity and addiction-like behaviour

Agent	Impulsive action		Impulsive choice		
	5-CSRTT	SSRTT/Go/No-go	DD	Self-administration	Reinstatement
Glutamate					
Non-competitive NMDA antagonist					
MK-801	↑ (Paine <i>et al.</i> , 2007)			↓FR, ↑PR cocaine (Allen <i>et al.</i> , 2005)	↓ cue-induced ethanol (von der Goltz <i>et al.</i> , 2009)
Ketamine	↑ (Oliver <i>et al.</i> , 2009) = (Nemeth <i>et al.</i> , 2010)		↑ (Cottone <i>et al.</i> , 2013) ↑ (Floresco <i>et al.</i> , 2008)	↓ FR ethanol (Sabino <i>et al.</i> , 2013)	
Memantine			↑ (Cottone <i>et al.</i> , 2013) = (Oberlin <i>et al.</i> , 2010)	↓ FR ethanol (Sabino <i>et al.</i> , 2013) ↓FR, PR cocaine (Hyytia <i>et al.</i> , 1999)	
PCP	↑ (Amitai <i>et al.</i> , 2007)			↓FR ethanol (Shelton and Balster, 1997)	
Selective glutamate receptor antagonist					
Ro63-1908 (NMDA 2B)	↑ (Higgins <i>et al.</i> , 2005; Burton and Fletcher, 2012)				
Ifenprodil (NMDA 2B)					↓ drug-primed ethanol (Vengeliene <i>et al.</i> , 2005) ↓ cue-induced nicotine (Gipson <i>et al.</i> , 2013)
EMQMCM (mGlu ₁)			↓ (Sukhotina <i>et al.</i> , 2008)		↓ cue-induced nicotine, drug-primed nicotine (Dravolina <i>et al.</i> , 2007)
JNJ16259685 (mGlu ₁)					↓ context- induced cocaine (Xie <i>et al.</i> , 2012b)
LY341495 (mGlu _{2/3})	= (Semenova and Markou, 2007)			↓FR, ↓PR cocaine (Allen <i>et al.</i> , 2005) ↓ FR ethanol (Backstrom and Hyytia, 2005)	↓ cue-induced ethanol (Backstrom and Hyytia, 2005) ↓ cue-induced cocaine (Baptista <i>et al.</i> , 2004)

Table 6

Continued

Agent	Impulsive action		Impulsive choice		
	5-CSRTT	SSRTT/Go/No-go	DD	Self-administration	Reinstatement
MPEP (mGlu ₅)	↓ (Semenova and Markou, 2007)			↓ FR ethanol (Schroeder <i>et al.</i> , 2005) ↓ FR, ↓ PR nicotine (Paterson and Markou, 2005) ↓ FR, PR cocaine (Paterson and Markou, 2005)	↓ cue-induced cocaine (Backstrom and Hyttia, 2006) ↓ cue-induced nicotine (Palmatier <i>et al.</i> , 2008) ↓ drug-primed nicotine (Tessari <i>et al.</i> , 2004)
GABA					
GABA-mimetic					
Ethanol	↑ (Oliver <i>et al.</i> , 2009)		↑ (Olmstead <i>et al.</i> , 2006)	↓ FR ethanol (Shelton and Balster, 1997)	
Vigabatrin				↓ FR cocaine (Filip <i>et al.</i> , 2007)	↓ drug-primed cocaine (Filip <i>et al.</i> , 2007)
GABA _A agonist					
Diazepam	↑ (Oliver <i>et al.</i> , 2009) = § (Molander <i>et al.</i> , 2011)		↑ (Thiebot <i>et al.</i> , 1985)	↓ FR cocaine (Augier <i>et al.</i> , 2012)	
Chlordiazepoxide			↑ (Cardinal <i>et al.</i> , 2000)		
GABA _B agonist					
Baclofen	↑ ILC (Murphy <i>et al.</i> , 2012) ↑ MRN (Le <i>et al.</i> , 2008) ↓ STN (Baunez and Robbins, 1999)		↑ OFC (Zeeb <i>et al.</i> , 2010) ↑ vHC (Abela and Chudasama, 2014) ↑ NAc Core (Ghods-Sharifi and Floresco, 2010) = NAc Shell (Ghods-Sharifi and Floresco, 2010) ↓ STN (Winstanley <i>et al.</i> , 2005a)	↓ FR, PR cocaine (Brebner <i>et al.</i> , 2000) ↓ FR nicotine (Fattore <i>et al.</i> , 2002)	↓ drug-primed nicotine (Fattore <i>et al.</i> , 2009)
Opioid					
Non-selective antagonist					
Naloxone	= (Pattij <i>et al.</i> , 2009; Wiskerke <i>et al.</i> , 2011b) = * (nicotine) (Wiskerke <i>et al.</i> , 2012) ↓ * (amphetamine, GBR12909) (Wiskerke <i>et al.</i> , 2011b)	= (Pattij <i>et al.</i> , 2009)	= (Pattij <i>et al.</i> , 2009) = / = * (amphetamine) (Wiskerke <i>et al.</i> , 2011b)	↓ FR cocaine (Kiyatkin and Brown, 2003) ↓ FR nicotine (Ismayilova and Shoaib, 2010)	↓ drug-primed cannabinoid (Spano <i>et al.</i> , 2004)

Table 6

Continued

Agent	Impulsive action		Impulsive choice		
	5-CSRTT	SSRTT/Go/No-go	DD	Self-administration	Reinstatement
μ agonist					
Morphine	↑ (Pattij <i>et al.</i> , 2009)	= (Pattij <i>et al.</i> , 2009)	↑ (Pattij <i>et al.</i> , 2009)		
δ antagonist					
Naltrindol	=/= * (amphetamine) (Wiskerke <i>et al.</i> , 2011b)		=/= * (amphetamine) (Wiskerke <i>et al.</i> , 2011b)	= FR nicotine (Ismayilova and Shoaib, 2010) = FR cocaine (de Vries <i>et al.</i> , 1995)	↓ cue-induced ethanol (Ciccocioppo <i>et al.</i> , 2002; Marinelli <i>et al.</i> , 2009)
κ agonist					
Salvinorin A	= (Nemeth <i>et al.</i> , 2010)				
U50 488		↑ (Walker and Kissler, 2013)	= (Walker and Kissler, 2013)	↓ FR nicotine (Ismayilova and Shoaib, 2010) ↓ FR cocaine (Glick <i>et al.</i> , 1995)	
U69593					↓ drug-primed cocaine (Schenk <i>et al.</i> , 2000)
κ antagonist					
Nor-BNI	=/= * (amphetamine, nicotine) (Wiskerke <i>et al.</i> , 2011b; 2012)	= (Walker and Kissler, 2013)	=/= * (amphetamine) (Wiskerke <i>et al.</i> , 2011b)	= FR cocaine (Glick <i>et al.</i> , 1995) ↓ PR cocaine (Wee <i>et al.</i> , 2009) ↓ FR ethanol (Walker <i>et al.</i> , 2011)	↓ stress- induced cocaine (Redila and Chavkin, 2008)
Cannabinoid					
CB ₁ agonist					
WIN 55212-2	= (Pattij <i>et al.</i> , 2007b)		= (Pattij <i>et al.</i> , 2007b)	↑ FR nicotine (Gamaledin <i>et al.</i> , 2012)	↑ drug-primed, cue- induced nicotine (Gamaledin <i>et al.</i> , 2012) ↑ drug-primed ethanol (Alen <i>et al.</i> , 2008)
CB ₁ antagonist					
SR141716A	=/= * (GBR12909)/↓* (nicotine) (Wiskerke <i>et al.</i> , 2012) ↓* (amphetamine) (Wiskerke <i>et al.</i> , 2011a) ↓ (Pattij <i>et al.</i> , 2007b)		= (Pattij <i>et al.</i> , 2007b) ↑* (amphetamine) (Wiskerke <i>et al.</i> , 2011a)	↓ FR, PR ethanol (Economidou <i>et al.</i> , 2006) = FR cocaine (De Vries <i>et al.</i> , 2001)	↓ cue-induced ethanol (Economidou <i>et al.</i> , 2006) ↓ cue-induced cocaine (De Vries <i>et al.</i> , 2001)

Table 6

Continued

Agent	Impulsive action		Impulsive choice		
	5-CSRTT	SSRTT/Go/No-go	DD	Self-administration	Reinstatement
SLV330	↓ (de Bruin <i>et al.</i> , 2011)			↓FR ethanol, nicotine (de Bruin <i>et al.</i> , 2011)	↓ cue-induced ethanol, nicotine (de Bruin <i>et al.</i> , 2011)
O-2050	↓/↓* (amphetamine) (Wiskerke <i>et al.</i> , 2011a)		↑* (amphetamine) (Wiskerke <i>et al.</i> , 2011a)		
Cholinergic nACh agonist					
Nicotine	↑ (Blondel <i>et al.</i> , 2000)	↑ (Kolokotroni <i>et al.</i> , 2011)	↑ (Dallery and Locey, 2005; Kolokotroni <i>et al.</i> , 2011)		
nACh antagonist					
Mecamylamine (α4β2)	↓ (Ruotsalainen <i>et al.</i> , 2000; Tsutsui-Kimura <i>et al.</i> , 2010) ↓* (nicotine) (Kolokotroni <i>et al.</i> , 2011)		= (Mendez <i>et al.</i> , 2012) ↓* (nicotine) (Kolokotroni <i>et al.</i> , 2011)	↓ FR cocaine (Levin <i>et al.</i> , 2000) ↓ FR ethanol (Kuzmin <i>et al.</i> , 2009) ↓ FR nicotine (Watkins <i>et al.</i> , 1999)	↓ cue-induced nicotine (Liu <i>et al.</i> , 2007)
Methyllaconitine (α7)	= (Tsutsui-Kimura <i>et al.</i> , 2010)			= FR ethanol (Le <i>et al.</i> , 2000)	= cue-induced nicotine (Liu, 2014) = drug-primed ethanol (Kuzmin <i>et al.</i> , 2009)
Non-selective mACh agonist					
Oxotremorine	= (Mirza and Stoleran, 2000)		= (Mendez <i>et al.</i> , 2012)	↓ FR cocaine (Rasmussen <i>et al.</i> , 2000; Thomsen <i>et al.</i> , 2010)	
Non-selective mACh antagonist					
Scopolamine	= (Ruotsalainen <i>et al.</i> , 2000) ↑ (Shannon and Eberle, 2006) ↓ (Mirza and Stoleran, 2000)		↑ (Mendez <i>et al.</i> , 2012)	↓ ethanol (Rezvani <i>et al.</i> , 1991)	↓ drug-primed cocaine (Yee <i>et al.</i> , 2011)
Atropine			↑ (Mendez <i>et al.</i> , 2012)	↓ FR amphetamine (Davis and Smith, 1975)	

*Denotes effect on pharmacologically increased levels of impulsivity, agent in parentheses.

§Denotes effect in selected high-impulsive rats.

↑, increased; ↓, decreased; =, no effect; BLA, basolateral amygdala; ILC, infralimbic cortex; MA, methamphetamine; mACh, muscarinic ACh receptor; MRN, median raphe nucleus; nACh, nicotinic ACh receptor; PCP, phencyclidine; STN, subthalamic nucleus; vHC, ventral hippocampus.

relapse to, many classes of abused drugs (e.g. Backstrom and Hyttia, 2005; Dravolina *et al.*, 2007; Xie *et al.*, 2010; 2012b).

In view of the predominantly consistent effect of antagonism at NMDA and mGlu receptors to enhance measures of action impulsivity, but *reduce* self-administration and relapse-like behaviour, it is unlikely that the link between action impulsivity and addiction is driven via activity in the glutamatergic system. There is the possibility, however, that action at mGlu₁ receptors may link addiction-related behaviours and impulsive choice given that antagonism at this receptor subtype commonly reduces both behavioural categories (e.g. Dravolina *et al.*, 2007; Sukhotina *et al.*, 2008).

GABAergic agents

While few studies have investigated the role of GABA in impulsivity (Hayes *et al.*, 2014), GABA_A and GABA_B agonists have generally been found to increase measures of impulsive action (Oliver *et al.*, 2009) and impulsive choice (Thiebot *et al.*, 1985; Cardinal *et al.*, 2000; Olmstead *et al.*, 2006; Table 6). However, enhancing GABAergic activity tends to decrease drug self-administration (Augier *et al.*, 2012) and relapse to drug-seeking (Filip *et al.*, 2007; Fattore *et al.*, 2009). Nevertheless, intracerebral infusions of GABA agonists have been shown to reduce impulsivity (e.g. Baunez and Robbins, 1999) suggesting that activity in local GABAergic microcircuits may bear a closer correspondence with impulsivity and addiction-related behaviours.

Opioidergic agents

Systemic administration of the non-selective μ -opioid receptor agonist morphine has been found to increase impulsivity in both delay discounting and the 5-CSRTT (Pattij *et al.*, 2009). At least for impulsive action, phasic activation at δ -opioid receptors has also been implicated in enhancing impulsivity (Befort *et al.*, 2011). Interestingly, antagonism at μ -opioid receptors has been shown to attenuate the effects of amphetamine and the dopamine re-uptake inhibitor GBR12909 to increase impulsivity in this task (Wiskerke *et al.*, 2011b), suggesting again that dopamine transmission is subject to modulation by a myriad of neurotransmitters, putatively at the level of the mesolimbic dopamine system (Diergaarde *et al.*, 2008). There is little evidence, however, for tonic activity at opioid receptors in mediating impulsive action or choice (e.g. Pattij *et al.*, 2009; Wiskerke *et al.*, 2011b; 2012).

Available evidence suggests some overlap between opioidergic mechanisms capable of affecting both impulsivity and addiction-related behaviours (Table 6). In general, μ - and δ -opioid receptor agonists are capable of enhancing drug self-administration (e.g. Sabino *et al.*, 2007) and relapse-like behaviour (e.g. Simmons and Self, 2009), although it should be noted that there was regional specificity in these effects (reviewed in Le Merrer *et al.*, 2009). Unlike the null findings for impulsivity, however, μ - and δ -opioid receptor antagonists generally reduce these behaviours [(e.g. Corrigan and Coen, 1991b; Ciccocioppo *et al.*, 2002; Kiyatkin and Brown, 2003; Spano *et al.*, 2004); for review, see van Ree *et al.*, 1999]. It remains to be seen whether such antagonists are capable of reducing impulsivity in animals with endogenously enhanced levels of this trait.

Cannabinoids

Despite a relative paucity of studies, the cannabinoid system offers potential scope for pharmacological intervention in both impulsivity and addiction. For example, tonic activity at cannabinoid type 1 receptors has been found to modulate nicotine-induced increases in impulsive responding on the 5-CSRTT (Wiskerke *et al.*, 2012). Furthermore, selective CB₁ receptor antagonists are capable of reducing baseline impulsivity as measured on this task (Pattij *et al.*, 2007b; de Bruin *et al.*, 2011); however, they have been found to have no effect on delay-discounting performance (Pattij *et al.*, 2007b). Systemic administration of CB₁ receptor antagonist SR141716A (rimonabant) in rats has been found to suppresses the self-administration of many drugs of abuse (reviewed in Maldonado *et al.*, 2006), including ethanol (e.g. Cippitelli *et al.*, 2005; Economidou *et al.*, 2006) and nicotine (e.g. Cohen *et al.*, 2002); however, findings for cocaine are variable (e.g. De Vries *et al.*, 2001; Soria *et al.*, 2005). The same compound also inhibits cue-induced cocaine, nicotine and ethanol reinstatement (e.g. De Vries *et al.*, 2001; 2005; Economidou *et al.*, 2006).

Cholinergic agents

A role for cholinergic mechanisms in impulsivity is supported by the common effect of nicotine to increase impulsive action (Blondel *et al.*, 2000; Kolokotroni *et al.*, 2011) and impulsive choice (Dallery and Locey, 2005; Kolokotroni *et al.*, 2011). Recent studies indicate that these effects may be mediated by nicotinic $\alpha 4 \beta 2$ receptors (Tsutsui-Kimura *et al.*, 2010; Xie *et al.*, 2012a). Interestingly, nicotinic $\alpha 4 \beta 2$ receptor antagonists also attenuate self-administration and relapse-like behaviour to nicotine and alcohol (Watkins *et al.*, 1999; Liu *et al.*, 2007; Kuzmin *et al.*, 2009). Additionally, activity at muscarinic cholinergic receptors has been found to modulate both measures of impulsivity and addiction-related behaviours. Although phasic muscarinic receptor activation appears to have no effect on impulsivity (Mirza and Stoleran, 2000; Mendez *et al.*, 2012), administration of non-selective muscarinic antagonists are reported to enhance delay discounting (Mendez *et al.*, 2012). However their effect on impulsive action is less clear with evidence for both increased (Shannon and Eberle, 2006) and decreased (Mirza and Stoleran, 2000) impulsivity on the 5-CSRTT. It is possible that variability of these findings may occur as a result of competing effects on attention (e.g. Ruotsalainen *et al.*, 2000). Other measures of behavioural inhibition such as DRL-72 and reaction time tasks, however, are generally impaired with reports of increased impulsivity (Blokland *et al.*, 2001; Jayarajan *et al.*, 2013). Interestingly, muscarinic M₁ receptor knock-out mice exhibit enhanced impulsivity on the 5-CSRTT independent of any effect on attention (Bartko *et al.*, 2011). There have also been discrepancies with the effect of muscarinic agents on drug reward and reinforcement. Non-selective muscarinic receptor agonists have been shown to reduce cocaine self-administration (Rasmussen *et al.*, 2000; Thomsen *et al.*, 2010), putatively mediated by M₁/M₄ receptors (Dencker *et al.*, 2012; Thomsen *et al.*, 2014). Interestingly, non-selective antagonism at muscarinic receptors has also been shown to reduce self-administration of methamphetamine (Davis and Smith, 1975) suggesting there may be drug and/or

receptor subtype-specific contributions of muscarinic signalling to drug reward and reinforcement.

Other neurotransmitter systems

Beyond those reviewed earlier, there is evidence for a role of other neurotransmitter systems in impulsivity and addiction-like behaviours. Despite being widely implicated in addiction (reviewed in Boutrel, 2008; Schank *et al.*, 2012), the contribution of stress-related neuropeptides to impulsive behaviour has been less well investigated. There is evidence that neurokinin 1 (NK₁) receptor antagonists reduce delay discounting (Loiseau *et al.*, 2005) as well as reduce stress-induced reinstatement of cocaine (Schank *et al.*, 2014) and alcohol (Schank *et al.*, 2011; 2014) seeking. Interestingly, however, antagonism at this receptor had no effect on cocaine (Placenza *et al.*, 2006) or alcohol reinforcement (Schank *et al.*, 2014). Neuropeptide Y, via an action at Y₂ receptors, has been reported to regulate impulsivity. Thus, Y₂ receptor knock-out mice were found to show increased impulsivity on the 5-CSRTT compared with wild-type littermates (Greco and Carli, 2006). To date, however, no studies have specifically investigated the effect of pharmacological manipulation of this system on measures of impulsivity in rodents. However, antagonism at Y₂ receptors has been shown to reduce responding for alcohol (Thorsell *et al.*, 2002; Rimondini *et al.*, 2005) although there are conflicting reports regarding this effect (Cippitelli *et al.*, 2011). Despite evidence for a role of corticotrophin releasing factor (CRF) signalling in addiction-related behaviours (reviewed in Logrip *et al.*, 2011) there is no evidence to suggest that CRF regulates impulsivity (Ohmura *et al.*, 2009).

Several other neurotransmitter systems that putatively interact with corticostriatal neurotransmission have also been implicated in both impulsivity and addiction. In this context, histamine H₃ receptors have been shown to form functional heterodimers with both D₁ and D₂ receptors (reviewed in Ellenbroek and Ghiabi, 2014). Antagonism at histamine H₃ receptors has been reported to reduce premature responding on the 5-CSRTT (Day *et al.*, 2007) and alcohol self-administration (Lintunen *et al.*, 2001; Nuutinen *et al.*, 2011), but enhance methamphetamine self-administration (Munzar *et al.*, 2004). Similarly, adenosine A₁ receptors form heteromeric complexes within the striatum to affect neurotransmission in this region (e.g. Ciruela *et al.*, 2006). Activation of this receptor subtype has been shown to reduce impulsive responding as measured on the DRL task (Marek, 2012). Additionally, activation of this receptor inhibits dopamine-induced relapse to cocaine-seeking (Hobson *et al.*, 2013), although its effects on other drug classes has yet to be investigated.

Clinical implications

Our analysis has revealed several promising pharmacological mechanisms that bisect impulsivity and addiction-related behaviours. As drug use in impulsive individuals may represent a form of self-mediation (Khantzian, 1985) treating co-morbid impulsive symptoms may help to curb continuing

drug use in addicts. While no studies to date have specifically investigated this hypothesis insights can perhaps be drawn from studies in ADHD.

With this in mind, a recent prospective study in adolescents diagnosed with ADHD and treated with methylphenidate found that rates of tobacco use were reduced when compared with a historical sample of smoking rates in a comparable, non-medicated population of individuals with ADHD (Hammerness *et al.*, 2013). However, a multi-site trial of the effect of methylphenidate treatment on rates of abstinence in individuals with ADHD and co-morbid tobacco dependence found no effect despite a reduction in ADHD symptoms (Covey *et al.*, 2011). These results plausibly suggest that treating impulsivity, at least with stimulant-based medications, may delay the development of addiction rather than remediating the active disease state once established. In keeping with this interpretation, meta-analyses suggest that stimulants protect against the development of later substance use disorders in individuals with ADHD (Wilens *et al.*, 2003), and that onset of treatment strongly predicts clinical outcome, with early initiation of treatment reducing the rate of dependence (Dalsgaard *et al.*, 2014). Clearly, much further research is needed to investigate the generality of these findings to all classes of abused drugs and whether NARIs (e.g. atomoxetine and roboxetine) offer similar protection against the development of addiction. In this regard, however, recent studies have found limited evidence for the efficacy of these drugs to enhance rates of abstinence in cocaine-dependent individuals with (Levin *et al.*, 2009) and without ADHD (Szerman *et al.*, 2005; Walsh *et al.*, 2013).

In light of enhanced rates of addiction, even in medicated ADHD patients (Hammerness *et al.*, 2013), there is naturally some concern whether exposing children and young adults to stimulant-based medications for ADHD might speed the development of later addiction. Preclinical studies have generally found that chronic pre-exposure to stimulant drugs increase subsequent levels of drug self-administration, at least in animals showing increased basal levels of impulsivity [i.e. spontaneous hypertensive rats, (Harvey *et al.*, 2011; Somkuwar *et al.*, 2013), but see Martelle *et al.* (2013); Gill *et al.* (2012); Thanos *et al.* (2007)]. However, more research is needed to understand the precise relationship between pre- and post-drug levels of impulsivity and whether in young animals stimulant exposure facilitates the development of compulsive drug self-administration, as previously described and modelled in rats (Vanderschuren and Everitt, 2004; Belin *et al.*, 2008; Pelloux *et al.*, 2012). In this regard, the reported action of atomoxetine to arrest the emergence of at least one form of compulsive behaviour in naturally impulsive rats (Ansquer *et al.*, 2014) strongly encourages further research in this area.

Conclusions

A review of the current literature highlights a number of convergent pharmacological mechanisms, which putatively mediate the reported link between impulsivity and addiction (see Table 7 for a summary). Interestingly, the available evidence suggests that while drugs capable of reducing

impulsive action often also decrease measures of addiction-related behaviours there is little convergence and often opposing effects with impulsive choice. The close alignment of pharmacological mechanisms in impulsive action and various animal addiction models suggests that they may tap into a common underlying process. Speculatively, this may involve modulatory effects on the invigoration of behaviour affecting the output of responses conditioned to food (impulsivity tasks) and drug (addiction tasks) rewards. This notion might then suggest a common involvement of mesolimbic dopamine projections, for example, to the ventral striatum, including the NAc, which are widely implicated in impulsivity (Basar *et al.*, 2010; Dalley *et al.*, 2011) and drug reinforcement (Willuhn *et al.*, 2010). Such interactions could operate at several levels including an enhancement or attenuation of appetitive approach behaviour to primary as well as conditioned reinforcing stimuli (Everitt and Robbins, 2005). Thus, the mesoaccumbens dopamine system may form the final common pathway through which other transmitter systems (for example those illustrated in Table 7) operate to modulate action impulsivity and certain drug-motivated behaviours (for reviews, see Feltenstein and See, 2013; Schmidt and Weinshenker, 2014). Conceivably, this mechanism may encompass dopamine-mediated gating of cortico-accumbens projections from the PFC, amygdala, and hippocampus (Goto and Grace, 2008), and thereby the expression of drug craving and relapse (Kalivas and Volkow, 2011). Interestingly, while our analysis has revealed a close relationship between motoric forms of impulsivity and addiction-related behaviours, there is little evidence for a similar association with impulsive choice, with the exception of nicotinic $\alpha 4\beta 2$ receptor antagonists, which decrease both action- and choice-related forms of impulsivity and drug-seeking responses. The apparently weak association between choice impulsivity and

addiction-related behaviours suggest they may involve distinct pharmacological substrates. Nevertheless, there is some evidence that NARIs may be efficacious in several forms of impulsivity, including delay-discounting impulsivity; however, larger-scale prospective studies are needed to investigate whether early intervention in ADHD (e.g. with NARIs) significantly moderates the risk of addiction in adulthood. Finally, research is needed to explore emerging neurotransmitter substrates and mechanisms (e.g. GABA, glutamate, cannabinoids and ACh) as a strategy to develop more effective therapies in addiction and impulsivity disorders.

Acknowledgements

This work was supported by the Medical Research Council (MRC G0701500) and by a joint award from the MRC and Wellcome Trust in support of the Behavioural and Clinical Neuroscience Institute at Cambridge University. The authors also acknowledge funding from the MRC Imperial College-Cambridge University-Manchester (ICCAM) strategic addiction cluster. BJ is supported by grants from the AXA Research Fund and the Australian National Health and Medical Research Council (1016313).

Author contributions

B. J. developed the review concept, wrote and edited the paper J. W. D. edited the paper, provided feedback during the drafting process and assisted in developing the review concept.

Table 7

Pharmacological interventions that reduce both impulsivity and addiction-like behaviour in rodent models

Neurotransmitter system	Impulsive action	Impulsive choice
Dopaminergic	D ₁ antagonist	
5-hydroxytryptaminergic	SSRI	
	5-HT _{2A} antagonist*	
	5-HT _{2A/C} antagonist	
	5-HT _{2C} agonist	
	5-HT _{2B/C} agonist	
	5-HT ₃ antagonist	
Adrenergic	NARI	
	NSRI*	
	α_1 antagonist	
	β antagonist	
Glutamate		mGlu ₁ antagonist
Cannabinoid	CB ₁ antagonist	
Cholinergic	Nicotinic $\alpha 4\beta 2$ antagonist	Nicotinic $\alpha 4\beta 2$ antagonist

*Evidence to suggest effect in relapse-like behaviour, but not self-administration, despite the apparent relationship of both to impulsivity. NSRI, noradrenaline 5-HT re-uptake inhibitor; SSRI, selective 5-HT re-uptake inhibitor.

Conflict of interest

None.

References

- Abela AR, Chudasama Y (2014). Noradrenergic alpha2A-receptor stimulation in the ventral hippocampus reduces impulsive decision-making. *Psychopharmacology (Berl)* 231: 521–531.
- Acosta JJ, Boynton FA, Kirschner KF, Neisewander JL (2005). Stimulation of 5-HT1B receptors decreases cocaine- and sucrose-seeking behavior. *Pharmacol Biochem Behav* 80: 297–307.
- Adriani W, Canese R, Podo F, Laviola G (2007). 1H MRS-detectable metabolic brain changes and reduced impulsive behavior in adult rats exposed to methylphenidate during adolescence. *Neurotoxicol Teratol* 29: 116–125.
- Agnoli L, Carli M (2011). Synergistic interaction of dopamine D(1) and glutamate N-methyl-D-aspartate receptors in the rat dorsal striatum controls attention. *Neuroscience* 185: 39–49.
- Agnoli L, Carli M (2012). Dorsal-striatal 5-HT(2)A and 5-HT(2)C receptors control impulsivity and perseverative responding in the 5-choice serial reaction time task. *Psychopharmacology (Berl)* 219: 633–645.
- Alen F, Moreno-Sanz G, Isabel de Tena A, Brooks RD, Lopez-Jimenez A, Navarro M *et al.* (2008). Pharmacological activation of CB1 and D2 receptors in rats: predominant role of CB1 in the increase of alcohol relapse. *Eur J Neurosci* 27: 3292–3298.
- Alexander SPH, Benson HE, Faccenda E, Pawson AJ, Sharman JL, Spedding M *et al.*; CGTP Collaborators (2013a). The Concise Guide to PHARMACOLOGY 2013/14: G protein-coupled receptors. *Br J Pharmacol* 170: 1459–1562.
- Alexander SPH, Benson HE, Faccenda E, Pawson AJ, Sharman JL, Spedding M *et al.*; CGTP Collaborators (2013b). The Concise Guide to PHARMACOLOGY 2013/14: Ligand-gated ion channels. *Br J Pharmacol* 170: 1582–1603.
- Allen RM, Carelli RM, Dykstra LA, Suchey TL, Everett CV (2005). Effects of the competitive N-methyl-D-aspartate receptor antagonist, LY235959 [(–)-6-phosphonomethyl-deca-hydroisoquinoline-3-carboxylic acid], on responding for cocaine under both fixed and progressive ratio schedules of reinforcement. *J Pharmacol Exp Ther* 315: 449–457.
- Amitai N, Semenova S, Markou A (2007). Cognitive-disruptive effects of the psychotomimetic phencyclidine and attenuation by atypical antipsychotic medications in rats. *Psychopharmacology (Berl)* 193: 521–537.
- Anderson SM, Schmidt HD, Pierce RC (2006). Administration of the D2 dopamine receptor antagonist sulpiride into the shell, but not the core, of the nucleus accumbens attenuates cocaine priming-induced reinstatement of drug seeking. *Neuropsychopharmacology* 31: 1452–1461.
- Ansquer S, Belin-Rauscent A, Dugast E, Duran T, Benatru I, Mar AC *et al.* (2014). Atomoxetine decreases vulnerability to develop compulsivity in high impulsive rats. *Biol Psychiatry* 75: 825–832.
- Augier E, Vouillac C, Ahmed SH (2012). Diazepam promotes choice of abstinence in cocaine self-administering rats. *Addict Biol* 17: 378–391.
- Avila C, Cuenca I, Felix V, Parcet MA, Miranda A (2004). Measuring impulsivity in school-aged boys and examining its relationship with ADHD and ODD ratings. *J Abnorm Child Psychol* 32: 295–304.
- Baarendse PJ, Vanderschuren LJ (2012). Dissociable effects of monoamine reuptake inhibitors on distinct forms of impulsive behavior in rats. *Psychopharmacology (Berl)* 219: 313–326.
- Bachtell RK, Whisler K, Karanian D, Self DW (2005). Effects of intra-nucleus accumbens shell administration of dopamine agonists and antagonists on cocaine-taking and cocaine-seeking behaviors in the rat. *Psychopharmacology (Berl)* 183: 41–53.
- Backstrom P, Hyytia P (2005). Suppression of alcohol self-administration and cue-induced reinstatement of alcohol seeking by the mGlu2/3 receptor agonist LY379268 and the mGlu8 receptor agonist (S)-3,4-DCPG. *Eur J Pharmacol* 528: 110–118.
- Backstrom P, Hyytia P (2006). Ionotropic and metabotropic glutamate receptor antagonism attenuates cue-induced cocaine seeking. *Neuropsychopharmacology* 31: 778–786.
- Banna KM, Back SE, Do P, See RE (2010). Yohimbine stress potentiates conditioned cue-induced reinstatement of heroin-seeking in rats. *Behav Brain Res* 208: 144–148.
- Baptista MA, Martin-Fardon R, Weiss F (2004). Preferential effects of the metabotropic glutamate 2/3 receptor agonist LY379268 on conditioned reinstatement versus primary reinforcement: comparison between cocaine and a potent conventional reinforcer. *J Neurosci* 24: 4723–4727.
- Barbelivien A, Billy E, Lazarus C, Kelche C, Majchrzak M (2008). Rats with different profiles of impulsive choice behavior exhibit differences in responses to caffeine and D-amphetamine and in medial prefrontal cortex 5-HT utilization. *Behav Brain Res* 187: 273–283.
- Bari A, Robbins TW (2013). Noradrenergic versus dopaminergic modulation of impulsivity, attention and monitoring behaviour in rats performing the stop-signal task: possible relevance to ADHD. *Psychopharmacology (Berl)* 230: 89–111.
- Bari A, Eagle DM, Mar AC, Robinson ES, Robbins TW (2009). Dissociable effects of noradrenaline, dopamine, and serotonin uptake blockade on stop task performance in rats. *Psychopharmacology (Berl)* 205: 273–283.
- Bari AA, Pierce RC (2005). D1-like and D2 dopamine receptor antagonists administered into the shell subregion of the rat nucleus accumbens decrease cocaine, but not food, reinforcement. *Neuroscience* 135: 959–968.
- Barkley RA, Fischer M, Smallish L, Fletcher K (2003). Does the treatment of attention-deficit/hyperactivity disorder with stimulants contribute to drug use/abuse? A 13-year prospective study. *Pediatrics* 111: 97–109.
- Barratt ES (1985) Impulsiveness subtraits, arousal and information processing. In: Spence JT, Itard CT (eds). *Motivation, Emotion and Personality*. Elsevier Science.: North Holland, pp. 137–146.
- Barrett AC, Miller JR, Dohrmann JM, Caine SB (2004). Effects of dopamine indirect agonists and selective D1-like and D2-like agonists and antagonists on cocaine self-administration and food maintained responding in rats. *Neuropharmacology* 47 (Suppl. 1): 256–273.
- Bartko SJ, Romberg C, White B, Wess J, Bussey TJ, Saksida LM (2011). Intact attentional processing but abnormal responding in M1 muscarinic receptor-deficient mice using an automated touchscreen method. *Neuropharmacology* 61: 1366–1378.

- Basar K, Sesia T, Groenewegen H, Steinbusch HW, Visser-Vandewalle V, Temel Y (2010). Nucleus accumbens and impulsivity. *Prog Neurobiol* 92: 533–557.
- Baunez C, Robbins TW (1999). Effects of transient inactivation of the subthalamic nucleus by local muscimol and APV infusions on performance on the five-choice serial reaction time task in rats. *Psychopharmacology (Berl)* 141: 57–65.
- Beftor K, Mahoney MK, Chow C, Hayton SJ, Kieffer BL, Olmstead MC (2011). Effects of delta opioid receptors activation on a response inhibition task in rats. *Psychopharmacology (Berl)* 214: 967–976.
- Belin D, Mar AC, Dalley JW, Robbins TW, Everitt BJ (2008). High impulsivity predicts the switch to compulsive cocaine-taking. *Science* 320: 1352–1355.
- van den Bergh FS, Bloemarts E, Groenink L, Olivier B, Oosting RS (2006). Delay aversion: effects of 7-OH-DPAT, 5-HT1A/1B-receptor stimulation and D-cycloserine. *Pharmacol Biochem Behav* 85: 736–743.
- Besson M, Belin D, McNamara R, Theobald DE, Castel A, Beckett VL *et al.* (2010). Dissociable control of impulsivity in rats by dopamine d2/3 receptors in the core and shell subregions of the nucleus accumbens. *Neuropsychopharmacology* 35: 560–569.
- Bizot J, Le Bihan C, Puech AJ, Hamon M, Thiebot M (1999). Serotonin and tolerance to delay of reward in rats. *Psychopharmacology (Berl)* 146: 400–412.
- Blasio A, Narayan AR, Kaminski BJ, Steardo L, Sabino V, Cottone P (2012). A modified adjusting delay task to assess impulsive choice between isocaloric reinforcers in non-deprived male rats: effects of 5-HT(2)A/C and 5-HT(1)A receptor agonists. *Psychopharmacology (Berl)* 219: 377–386.
- Blokhina EA, Kashkin VA, Zvartau EE, Danysz W, Bessalov AY (2005). Effects of nicotinic and NMDA receptor channel blockers on intravenous cocaine and nicotine self-administration in mice. *Eur Neuropsychopharmacol* 15: 219–225.
- Blokland A, Scholtissen B, Vermeeren A, Ramaekers J (2001). Dissociable effects of histamine H1 antagonists on reaction-time performance in rats. *Pharmacol Biochem Behav* 70: 427–436.
- Blokland A, Sik A, Lieben C (2005). Evaluation of DOI, 8-OH-DPAT, eticlopride and amphetamine on impulsive responding in a reaction time task in rats. *Behav Pharmacol* 16: 93–100.
- Blondeau C, Dellu-Hagedorn F (2007). Dimensional analysis of ADHD subtypes in rats. *Biol Psychiatry* 61: 1340–1350.
- Blondel A, Sanger DJ, Moser PC (2000). Characterisation of the effects of nicotine in the five-choice serial reaction time task in rats: antagonist studies. *Psychopharmacology (Berl)* 149: 293–305.
- Boutrel B (2008). A neuropeptide-centric view of psychostimulant addiction. *Br J Pharmacol* 154: 343–357.
- Brebner K, Phelan R, Roberts DC (2000). Effect of baclofen on cocaine self-administration in rats reinforced under fixed-ratio 1 and progressive-ratio schedules. *Psychopharmacology (Berl)* 148: 314–321.
- van den Brink W (2012). Evidence-based pharmacological treatment of substance use disorders and pathological gambling. *Curr Drug Abuse Rev* 5: 3–31.
- Broos N, Diergaarde L, Schoffelmeer AN, Pattij T, De Vries TJ (2012a). Trait impulsive choice predicts resistance to extinction and propensity to relapse to cocaine seeking: a bidirectional investigation. *Neuropsychopharmacology* 37: 1377–1386.
- Broos N, Schmaal L, Wiskerke J, Kostelijk L, Lam T, Stoop N *et al.* (2012b). The relationship between impulsive choice and impulsive action: a cross-species translational study. *PLoS ONE* 7: e36781.
- Bruijnzeel AW, Markou A (2003). Characterization of the effects of bupropion on the reinforcing properties of nicotine and food in rats. *Synapse* 50: 20–28.
- de Bruin NM, Lange JH, Kruse CG, Herremans AH, Schoffelmeer AN, van Drimmelen M *et al.* (2011). SLV330, a cannabinoid CB(1) receptor antagonist, attenuates ethanol and nicotine seeking and improves inhibitory response control in rats. *Behav Brain Res* 217: 408–415.
- de Bruin NM, McCreary AC, van Loevezijn A, de Vries TJ, Venhorst J, van Drimmelen M *et al.* (2013). A novel highly selective 5-HT6 receptor antagonist attenuates ethanol and nicotine seeking but does not affect inhibitory response control in Wistar rats. *Behav Brain Res* 236: 157–165.
- Buffalari DM, Baldwin CK, See RE (2012). Treatment of cocaine withdrawal anxiety with guanfacine: relationships to cocaine intake and reinstatement of cocaine seeking in rats. *Psychopharmacology (Berl)* 223: 179–190.
- Burmeister JJ, Lungren EM, Kirschner KF, Neisewander JL (2004). Differential roles of 5-HT receptor subtypes in cue and cocaine reinstatement of cocaine-seeking behavior in rats. *Neuropsychopharmacology* 29: 660–668.
- Burton CL, Fletcher PJ (2012). Age and sex differences in impulsive action in rats: the role of dopamine and glutamate. *Behav Brain Res* 230: 21–33.
- Caine SB, Koob GF (1993). Modulation of cocaine self-administration in the rat through D-3 dopamine receptors. *Science* 260: 1814–1816.
- Caine SB, Koob GF (1994). Effects of dopamine D-1 and D-2 antagonists on cocaine self-administration under different schedules of reinforcement in the rat. *J Pharmacol Exp Ther* 270: 209–218.
- Caine SB, Koob GF (1995). Pretreatment with the dopamine agonist 7-OH-DPAT shifts the cocaine self-administration dose-effect function to the left under different schedules in the rat. *Behav Pharmacol* 6: 333–347.
- Caine SB, Heinrichs SC, Coffin VL, Koob GF (1995). Effects of the dopamine D-1 antagonist SCH 23390 microinjected into the accumbens, amygdala or striatum on cocaine self-administration in the rat. *Brain Res* 692: 47–56.
- Caine SB, Negus SS, Mello NK, Bergman J (1999). Effects of dopamine D(1-like) and D(2-like) agonists in rats that self-administer cocaine. *J Pharmacol Exp Ther* 291: 353–360.
- Capriles N, Rodaros D, Sorge RE, Stewart J (2003). A role for the prefrontal cortex in stress- and cocaine-induced reinstatement of cocaine seeking in rats. *Psychopharmacology (Berl)* 168: 66–74.
- Caprioli D, Hong YT, Sawiak SJ, Ferrari V, Williamson DJ, Jupp B *et al.* (2013). Baseline-dependent effects of cocaine pre-exposure on impulsivity and D2/3 receptor availability in the rat striatum: possible relevance to the attention-deficit hyperactivity syndrome. *Neuropsychopharmacology* 38: 1460–1471.
- Carati C, Schenk S (2011). Role of dopamine D1- and D2-like receptor mechanisms in drug-seeking following methamphetamine self-administration in rats. *Pharmacol Biochem Behav* 98: 449–454.
- Cardinal RN (2006). Neural systems implicated in delayed and probabilistic reinforcement. *Neural Netw* 19: 1277–1301.
- Cardinal RN, Robbins TW, Everitt BJ (2000). The effects of d-amphetamine, chlordiazepoxide, alpha-flupenthixol and

behavioural manipulations on choice of signalled and unsignalled delayed reinforcement in rats. *Psychopharmacology (Berl)* 152: 362–375.

Carli M, Samanin R (2000). The 5-HT(1A) receptor agonist 8-OH-DPAT reduces rats' accuracy of attentional performance and enhances impulsive responding in a five-choice serial reaction time task: role of presynaptic 5-HT(1A) receptors. *Psychopharmacology (Berl)* 149: 259–268.

Carli M, Baviera M, Invernizzi RW, Balducci C (2006). Dissociable contribution of 5-HT1A and 5-HT2A receptors in the medial prefrontal cortex to different aspects of executive control such as impulsivity and compulsive perseveration in rats. *Neuropsychopharmacology* 31: 757–767.

Chakroun N, Doron J, Swendsen J (2004). Substance use, affective problems and personality traits: test of two association models]. *Encephale* 30: 564–569.

Chamberlain SR, Sahakian BJ (2007). The neuropsychiatry of impulsivity. *Curr Opin Psychiatry* 20: 255–261.

Chaudhri N, Sahuque LL, Janak PH (2009). Ethanol seeking triggered by environmental context is attenuated by blocking dopamine D1 receptors in the nucleus accumbens core and shell in rats. *Psychopharmacology (Berl)* 207: 303–314.

Chiamulera C, Tedesco V, Zangrandi L, Giuliano C, Fumagalli G (2010). Propranolol transiently inhibits reinstatement of nicotine-seeking behaviour in rats. *J Psychopharmacol* 24: 389–395.

Ciccocioppo R, Martin-Fardon R, Weiss F (2002). Effect of selective blockade of mu(1) or delta opioid receptors on reinstatement of alcohol-seeking behavior by drug-associated stimuli in rats. *Neuropsychopharmacology* 27: 391–399.

Cippitelli A, Bilbao A, Hansson AC, del Arco I, Sommer W, Heilig M *et al.* (2005). Cannabinoid CB1 receptor antagonism reduces conditioned reinstatement of ethanol-seeking behavior in rats. *Eur J Neurosci* 21: 2243–2251.

Cippitelli A, Rezvani AH, Robinson JE, Eisenberg L, Levin ED, Bonaventure P *et al.* (2011). The novel, selective, brain-penetrant neuropeptide Y Y2 receptor antagonist, JNJ-31020028, tested in animal models of alcohol consumption, relapse, and anxiety. *Alcohol* 45: 567–576.

Ciruela F, Casado V, Rodrigues RJ, Lujan R, Burgueno J, Canals M *et al.* (2006). Presynaptic control of striatal glutamatergic neurotransmission by adenosine A1-A2A receptor heteromers. *J Neurosci* 26: 2080–2087.

Cohen C, Perrault G, Sanger DJ (1998). Preferential involvement of D3 versus D2 dopamine receptors in the effects of dopamine receptor ligands on oral ethanol self-administration in rats. *Psychopharmacology (Berl)* 140: 478–485.

Cohen C, Perrault G, Sanger DJ (1999). Effects of D1 dopamine receptor agonists on oral ethanol self-administration in rats: comparison with their efficacy to produce grooming and hyperactivity. *Psychopharmacology (Berl)* 142: 102–110.

Cohen C, Perrault G, Voltz C, Steinberg R, Soubrie P (2002). SR141716, a central cannabinoid (CB1) receptor antagonist, blocks the motivational and dopamine-releasing effects of nicotine in rats. *Behav Pharmacol* 13: 451–463.

Cole BJ, Robbins TW (1987). Amphetamine impairs the discriminative performance of rats with dorsal noradrenergic bundle lesions on a 5-choice serial reaction time task: new evidence for central dopaminergic-noradrenergic interactions. *Psychopharmacology (Berl)* 91: 458–466.

Corrigall WA, Coen KM (1991a). Cocaine self-administration is increased by both D1 and D2 dopamine antagonists. *Pharmacol Biochem Behav* 39: 799–802.

Corrigall WA, Coen KM (1991b). Opiate antagonists reduce cocaine but not nicotine self-administration. *Psychopharmacology (Berl)* 104: 167–170.

Corrigall WA, Coen KM (1994). Nicotine self-administration and locomotor activity are not modified by the 5-HT3 antagonists ICS 205-930 and MDL 72222. *Pharmacol Biochem Behav* 49: 67–71.

Cottone P, Iemolo A, Narayan AR, Kwak J, Momany D, Sabino V (2013). The uncompetitive NMDA receptor antagonists ketamine and memantine preferentially increase the choice for a small, immediate reward in low-impulsive rats. *Psychopharmacology (Berl)* 226: 127–138.

Covey LS, Hu MC, Green CA, Brigham G, Hurt RD, Adler L *et al.* (2011). An exploration of site effects in a multisite trial of OROS-methylphenidate for smokers with attention deficit/hyperactivity disorder. *Am J Drug Alcohol Abuse* 37: 392–399.

Crombag HS, Grimm JW, Shaham Y (2002). Effect of dopamine receptor antagonists on renewal of cocaine seeking by reexposure to drug-associated contextual cues. *Neuropsychopharmacology* 27: 1006–1015.

Cunningham KA, Fox RG, Anastasio NC, Bubar MJ, Stutz SJ, Moeller FG *et al.* (2011). Selective serotonin 5-HT(2C) receptor activation suppresses the reinforcing efficacy of cocaine and sucrose but differentially affects the incentive-salience value of cocaine- vs. sucrose-associated cues. *Neuropharmacology* 61: 513–523.

Dallery J, Locey ML (2005). Effects of acute and chronic nicotine on impulsive choice in rats. *Behav Pharmacol* 16: 15–23.

Dalley JW, Roiser JP (2012). Dopamine, serotonin and impulsivity. *Neuroscience* 215: 42–58.

Dalley JW, Fryer TD, Brichard L, Robinson ES, Theobald DE, Laane K *et al.* (2007). Nucleus accumbens D2/3 receptors predict trait impulsivity and cocaine reinforcement. *Science* 315: 1267–1270.

Dalley JW, Everitt BJ, Robbins TW (2011). Impulsivity, compulsivity, and top-down cognitive control. *Neuron* 69: 680–694.

Dalsgaard S, Mortensen PB, Frydenberg M, Thomsen PH (2014). ADHD, stimulant treatment in childhood and subsequent substance abuse in adulthood – a naturalistic long-term follow-up study. *Addict Behav* 39: 325–328.

Daruma J, Barnes P (1993) A neurodevelopmental view of impulsivity and its relationship to the superfactors of personality. In: McCown W, Johnson J, Shure M (eds). *The Impulsive Client: Theory, Research and Treatment*. American Psychological Association: Washington DC, p. 23.

David V, Segu L, Buhot MC, Ichaye M, Cazala P (2004). Rewarding effects elicited by cocaine microinjections into the ventral tegmental area of C57BL/6 mice: involvement of dopamine D1 and serotonin1B receptors. *Psychopharmacology (Berl)* 174: 367–375.

David V, Besson M, Changeux JP, Granon S, Cazala P (2006). Reinforcing effects of nicotine microinjections into the ventral tegmental area of mice: dependence on cholinergic nicotinic and dopaminergic D1 receptors. *Neuropharmacology* 50: 1030–1040.

Davidson C, Lee TH, Xiong Z, Ellinwood EH (2002). Ondansetron given in the acute withdrawal from a repeated cocaine sensitization dosing regimen reverses the expression of sensitization and inhibits self-administration. *Neuropsychopharmacology* 27: 542–553.

Davis WM, Smith SG (1975). Central cholinergic influence on self-administration of morphine and amphetamine. *Life Sci* 16: 237–246.

- Day M, Pan JB, Buckley MJ, Cronin E, Hollingsworth PR, Hirst WD *et al.* (2007). Differential effects of ciprofloxacin and nicotine on impulsivity and attention measures in the 5-choice serial reaction time test. *Biochem Pharmacol* 73: 1123–1134.
- De Vries TJ, Schoffelmeier AN, Binnekade R, Mulder AH, Vanderschuren LJ (1998a). Drug-induced reinstatement of heroin- and cocaine-seeking behaviour following long-term extinction is associated with expression of behavioural sensitization. *Eur J Neurosci* 10: 3565–3571.
- De Vries TJ, Schoffelmeier AN, Binnekade R, Mulder AH, Vanderschuren LJ (1998b). MK-801 reinstates drug-seeking behaviour in cocaine-trained rats. *Neuroreport* 9: 637–640.
- De Vries TJ, Schoffelmeier AN, Binnekade R, Vanderschuren LJ (1999). Dopaminergic mechanisms mediating the incentive to seek cocaine and heroin following long-term withdrawal of IV drug self-administration. *Psychopharmacology (Berl)* 143: 254–260.
- De Vries TJ, Shaham Y, Homberg JR, Crombag H, Schuurman K, Dieben J *et al.* (2001). A cannabinoid mechanism in relapse to cocaine seeking. *Nat Med* 7: 1151–1154.
- De Vries TJ, Schoffelmeier AN, Binnekade R, Raaso H, Vanderschuren LJ (2002). Relapse to cocaine- and heroin-seeking behavior mediated by dopamine D2 receptors is time-dependent and associated with behavioral sensitization. *Neuropsychopharmacology* 26: 18–26.
- De Vries TJ, de Vries W, Janssen MC, Schoffelmeier AN (2005). Suppression of conditioned nicotine and sucrose seeking by the cannabinoid-1 receptor antagonist SR141716A. *Behav Brain Res* 161: 164–168.
- Del Campo N, Chamberlain SR, Sahakian BJ, Robbins TW (2011). The roles of dopamine and noradrenaline in the pathophysiology and treatment of attention-deficit/hyperactivity disorder. *Biol Psychiatry* 69: e145–e157.
- Dencker D, Weikop P, Sorensen G, Woldbye DP, Wortwein G, Wess J *et al.* (2012). An allosteric enhancer of M(4) muscarinic acetylcholine receptor function inhibits behavioral and neurochemical effects of cocaine. *Psychopharmacology (Berl)* 224: 277–287.
- Depoortere RY, Li DH, Lane JD, Emmett-Oglesby MW (1993). Parameters of self-administration of cocaine in rats under a progressive-ratio schedule. *Pharmacol Biochem Behav* 45: 539–548.
- Deroche-Gamonet V, Belin D, Piazza PV (2004). Evidence for addiction-like behavior in the rat. *Science* 305: 1014–1017.
- Di Chiara G, Imperato A (1988). Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci U S A* 85: 5274–5278.
- Di Matteo V, Di Giovanni G, Pierucci M, Esposito E (2008). Serotonin control of central dopaminergic function: focus on *in vivo* microdialysis studies. *Prog Brain Res* 172: 7–44.
- Dickman SJ (1990). Functional and dysfunctional impulsivity: personality and cognitive correlates. *J Pers Soc Psychol* 58: 95–102.
- Diergaarde L, Pattij T, Poortvliet I, Hogenboom F, de Vries W, Schoffelmeier AN *et al.* (2008). Impulsive choice and impulsive action predict vulnerability to distinct stages of nicotine seeking in rats. *Biol Psychiatry* 63: 301–308.
- Ding ZM, Toalston JE, Oster SM, McBride WJ, Rodd ZA (2009). Involvement of local serotonin-2A but not serotonin-1B receptors in the reinforcing effects of ethanol within the posterior ventral tegmental area of female Wistar rats. *Psychopharmacology (Berl)* 204: 381–390.
- Dougherty DM, Mathias CW, Marsh DM, Papageorgiou TD, Swann AC, Moeller FG (2004). Laboratory measured behavioral impulsivity relates to suicide attempt history. *Suicide Life Threat Behav* 34: 374–385.
- Dravolina OA, Zakharchova ES, Shekunova EV, Zvartau EE, Danysz W, Bessalov AY (2007). mGlu1 receptor blockade attenuates cue- and nicotine-induced reinstatement of extinguished nicotine self-administration behavior in rats. *Neuropharmacology* 52: 263–269.
- Dyr W, McBride WJ, Lumeng L, Li TK, Murphy JM (1993). Effects of D1 and D2 dopamine receptor agents on ethanol consumption in the high-alcohol-drinking (HAD) line of rats. *Alcohol* 10: 207–212.
- Eagle DM, Robbins TW (2003). Lesions of the medial prefrontal cortex or nucleus accumbens core do not impair inhibitory control in rats performing a stop-signal reaction time task. *Behav Brain Res* 146: 131–144.
- Eagle DM, Tufft MR, Goodchild HL, Robbins TW (2007). Differential effects of modafinil and methylphenidate on stop-signal reaction time task performance in the rat, and interactions with the dopamine receptor antagonist cis-flupenthixol. *Psychopharmacology (Berl)* 192: 193–206.
- Eagle DM, Lehmann O, Theobald DE, Pena Y, Zakaria R, Ghosh R *et al.* (2009). Serotonin depletion impairs waiting but not stop-signal reaction time in rats: implications for theories of the role of 5-HT in behavioral inhibition. *Neuropsychopharmacology* 34: 1311–1321.
- Eagle DM, Wong JC, Allan ME, Mar AC, Theobald DE, Robbins TW (2011). Contrasting roles for dopamine D1 and D2 receptor subtypes in the dorsomedial striatum but not the nucleus accumbens core during behavioral inhibition in the stop-signal task in rats. *J Neurosci* 31: 7349–7356.
- Ecke LE, Elmer GI, Suto N (2012). Cocaine self-administration is not dependent upon mesocortical alpha1 noradrenergic signaling. *Neuroreport* 23: 325–330.
- Economidou D, Mattioli L, Cifani C, Perfumi M, Massi M, Cuomo V *et al.* (2006). Effect of the cannabinoid CB1 receptor antagonist SR-141716A on ethanol self-administration and ethanol-seeking behaviour in rats. *Psychopharmacology (Berl)* 183: 394–403.
- Economidou D, Pelloux Y, Robbins TW, Dalley JW, Everitt BJ (2009). High impulsivity predicts relapse to cocaine-seeking after punishment-induced abstinence. *Biol Psychiatry* 65: 851–856.
- Economidou D, Dalley JW, Everitt BJ (2011). Selective norepinephrine reuptake inhibition by atomoxetine prevents cue-induced heroin and cocaine seeking. *Biol Psychiatry* 69: 266–274.
- Economidou D, Theobald DE, Robbins TW, Everitt BJ, Dalley JW (2012). Norepinephrine and dopamine modulate impulsivity on the five-choice serial reaction time task through opponent actions in the shell and core sub-regions of the nucleus accumbens. *Neuropsychopharmacology* 37: 2057–2066.
- Edwards S, Whisler KN, Fuller DC, Orsulak PJ, Self DW (2007). Addiction-related alterations in D1 and D2 dopamine receptor behavioral responses following chronic cocaine self-administration. *Neuropsychopharmacology* 32: 354–366.
- Ellenbroek BA, Ghiabi B (2014). The other side of the histamine H3 receptor. *Trends Neurosci* 37: 191–199.
- Erb S, Hitchcott PK, Rajabi H, Mueller D, Shaham Y, Stewart J (2000). Alpha-2 adrenergic receptor agonists block stress-induced reinstatement of cocaine seeking. *Neuropsychopharmacology* 23: 138–150.

- Ersche KD, Turton AJ, Pradhan S, Bullmore ET, Robbins TW (2010). Drug addiction endophenotypes: impulsive versus sensation-seeking personality traits. *Biol Psychiatry* 68: 770–773.
- Evernden JL (1999a). Varieties of impulsivity. *Psychopharmacology (Berl)* 146: 348–361.
- Evernden JL (1999b). The pharmacology of impulsive behaviour in rats VII: the effects of serotonergic agonists and antagonists on responding under a discrimination task using unreliable visual stimuli. *Psychopharmacology (Berl)* 146: 422–431.
- Evernden JL, Ryan CN (1996). The pharmacology of impulsive behaviour in rats: the effects of drugs on response choice with varying delays of reinforcement. *Psychopharmacology (Berl)* 128: 161–170.
- Everitt BJ, Robbins TW (2005). Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat Neurosci* 8: 1481–1489.
- Faraone SV, Biederman J, Spencer T, Michelson D, Adler L, Reimherr F *et al.* (2005). Efficacy of atomoxetine in adult attention-deficit/hyperactivity disorder: a drug-placebo response curve analysis. *Behav Brain Funct* 1: 16.
- Fattore L, Cossu G, Martellotta MC, Fratta W (2002). Baclofen antagonizes intravenous self-administration of nicotine in mice and rats. *Alcohol Alcohol* 37: 495–498.
- Fattore L, Spano MS, Cossu G, Scherma M, Fratta W, Fadda P (2009). Baclofen prevents drug-induced reinstatement of extinguished nicotine-seeking behaviour and nicotine place preference in rodents. *Eur Neuropsychopharmacol* 19: 487–498.
- Feltenstein MW, See RE (2006). Potentiation of cue-induced reinstatement of cocaine-seeking in rats by the anxiogenic drug yohimbine. *Behav Brain Res* 174: 1–8.
- Feltenstein MW, See RE (2013). Systems level neuroplasticity in drug addiction. *Cold Spring Harb Perspect Med* 3: a011916.
- Feltenstein MW, Altar CA, See RE (2007). Aripiprazole blocks reinstatement of cocaine seeking in an animal model of relapse. *Biol Psychiatry* 61: 582–590.
- Feltenstein MW, Henderson AR, See RE (2011). Enhancement of cue-induced reinstatement of cocaine-seeking in rats by yohimbine: sex differences and the role of the estrous cycle. *Psychopharmacology (Berl)* 216: 53–62.
- Feltenstein MW, Ghee SM, See RE (2012). Nicotine self-administration and reinstatement of nicotine-seeking in male and female rats. *Drug Alcohol Depend* 121: 240–246.
- Feola TW, de Wit H, Richards JB (2000). Effects of d-amphetamine and alcohol on a measure of behavioral inhibition in rats. *Behav Neurosci* 114: 838–848.
- Fernando AB, Economidou D, Theobald DE, Zou MF, Newman AH, Spoelder M *et al.* (2012). Modulation of high impulsivity and attentional performance in rats by selective direct and indirect dopaminergic and noradrenergic receptor agonists. *Psychopharmacology (Berl)* 219: 341–352.
- Filip M (2005). Role of serotonin (5-HT)₂ receptors in cocaine self-administration and seeking behavior in rats. *Pharmacol Rep* 57: 35–46.
- Filip M, Frankowska M, Zaniwska M, Golda A, Przegalinski E, Vetulani J (2007). Diverse effects of GABA-mimetic drugs on cocaine-evoked self-administration and discriminative stimulus effects in rats. *Psychopharmacology (Berl)* 192: 17–26.
- Filip M, Alenina N, Bader M, Przegalinski E (2010). Behavioral evidence for the significance of serotonergic (5-HT) receptors in cocaine addiction. *Addict Biol* 15: 227–249.
- Fineberg NA, Chamberlain SR, Goudriaan AE, Stein DJ, Vanderschuren LJ, Gillan CM *et al.* (2014). New developments in human neurocognition: clinical, genetic, and brain imaging correlates of impulsivity and compulsivity. *CNS Spectr* 19: 69–89.
- Fink KB, Gothert M (2007). 5-HT receptor regulation of neurotransmitter release. *Pharmacol Rev* 59: 360–417.
- Fletcher PJ (1998). A comparison of the effects of risperidone, raclopride, and ritanserin on intravenous self-administration of d-amphetamine. *Pharmacol Biochem Behav* 60: 55–60.
- Fletcher PJ, Korth KM, Chambers JW (1999). Depletion of brain serotonin following intra-raphe injections of 5,7-dihydroxytryptamine does not alter d-amphetamine self-administration across different schedule and access conditions. *Psychopharmacology (Berl)* 146: 185–193.
- Fletcher PJ, Grottick AJ, Higgins GA (2002). Differential effects of the 5-HT(2A) receptor antagonist M100907 and the 5-HT(2C) receptor antagonist SB242084 on cocaine-induced locomotor activity, cocaine self-administration and cocaine-induced reinstatement of responding. *Neuropsychopharmacology* 27: 576–586.
- Fletcher PJ, Chintoh AF, Sinyard J, Higgins GA (2004). Injection of the 5-HT_{2C} receptor agonist Ro60-0175 into the ventral tegmental area reduces cocaine-induced locomotor activity and cocaine self-administration. *Neuropsychopharmacology* 29: 308–318.
- Fletcher PJ, Tampakeras M, Sinyard J, Higgins GA (2007). Opposing effects of 5-HT(2A) and 5-HT(2C) receptor antagonists in the rat and mouse on premature responding in the five-choice serial reaction time test. *Psychopharmacology (Berl)* 195: 223–234.
- Fletcher PJ, Rizos Z, Sinyard J, Tampakeras M, Higgins GA (2008). The 5-HT_{2C} receptor agonist Ro60-0175 reduces cocaine self-administration and reinstatement induced by the stressor yohimbine, and contextual cues. *Neuropsychopharmacology* 33: 1402–1412.
- Fletcher PJ, Chambers JW, Rizos Z, Chintoh AF (2009). Effects of 5-HT depletion in the frontal cortex or nucleus accumbens on response inhibition measured in the 5-choice serial reaction time test and on a DRL schedule. *Behav Brain Res* 201: 88–98.
- Fletcher PJ, Rizos Z, Noble K, Higgins GA (2011). Impulsive action induced by amphetamine, cocaine and MK801 is reduced by 5-HT(2C) receptor stimulation and 5-HT(2A) receptor blockade. *Neuropharmacology* 61: 468–477.
- Fletcher PJ, Rizos Z, Noble K, Soko AD, Silenieks LB, Le AD *et al.* (2012). Effects of the 5-HT_{2C} receptor agonist Ro60-0175 and the 5-HT_{2A} receptor antagonist M100907 on nicotine self-administration and reinstatement. *Neuropharmacology* 62: 2288–2298.
- Floresco SB, Tse MT, Ghods-Sharifi S (2008). Dopaminergic and glutamatergic regulation of effort- and delay-based decision making. *Neuropsychopharmacology* 33: 1966–1979.
- Fone KC, Nutt DJ (2005). Stimulants: use and abuse in the treatment of attention deficit hyperactivity disorder. *Curr Opin Pharmacol* 5: 87–93.
- Forget B, Wertheim C, Mascia P, Pushparaj A, Goldberg SR, Le Foll B (2010). Noradrenergic α 1 receptors as a novel target for the treatment of nicotine addiction. *Neuropsychopharmacology* 35: 1751–1760.
- van Gaalen MM, van Koten R, Schoffelmeer AN, Vanderschuren LJ (2006a). Critical involvement of dopaminergic neurotransmission in impulsive decision making. *Biol Psychiatry* 60: 66–73.

- van Gaalen MM, Brueggeman RJ, Bronius PF, Schoffelman AN, Vanderschuren LJ (2006b). Behavioral disinhibition requires dopamine receptor activation. *Psychopharmacology (Berl)* 187: 73–85.
- van Gaalen MM, Unger L, Jongen-Relo AL, Schoemaker H, Gross G (2009). Amphetamine decreases behavioral inhibition by stimulation of dopamine D2, but not D3, receptors. *Behav Pharmacol* 20: 484–491.
- Gamaledin I, Wertheim C, Zhu AZ, Coen KM, Vemuri K, Makryannis A *et al.* (2012). Cannabinoid receptor stimulation increases motivation for nicotine and nicotine seeking. *Addict Biol* 17: 47–61.
- Garavan H, Kaufman JN, Hester R (2008). Acute effects of cocaine on the neurobiology of cognitive control. *Philos Trans R Soc Lond B Biol Sci* 363: 3267–3276.
- Gass JT, Olive MF (2008). Glutamatergic substrates of drug addiction and alcoholism. *Biochem Pharmacol* 75: 218–265.
- George O, Le Moal M, Koob GF (2012). Allostasis and addiction: role of the dopamine and corticotropin-releasing factor systems. *Physiol Behav* 106: 58–64.
- Ghods-Sharifi S, Floresco SB (2010). Differential effects on effort discounting induced by inactivations of the nucleus accumbens core or shell. *Behav Neurosci* 124: 179–191.
- Gill KE, Pierre PJ, Daunais J, Bennett AJ, Martelle S, Gage HD *et al.* (2012). Chronic treatment with extended release methylphenidate does not alter dopamine systems or increase vulnerability for cocaine self-administration: a study in nonhuman primates. *Neuropsychopharmacology* 37: 2555–2565.
- Gilpin NW, Koob GF (2010). Effects of beta-adrenoceptor antagonists on alcohol drinking by alcohol-dependent rats. *Psychopharmacology (Berl)* 212: 431–439.
- Gipson CD, Reissner KJ, Kupchik YM, Smith AC, Stankeviciute N, Hensley-Simon ME *et al.* (2013). Reinstatement of nicotine seeking is mediated by glutamatergic plasticity. *Proc Natl Acad Sci U S A* 110: 9124–9129.
- Glatz AC, Ehrlich M, Bae RS, Clarke MJ, Quinlan PA, Brown EC *et al.* (2002). Inhibition of cocaine self-administration by fluoxetine or D-fenfluramine combined with phentermine. *Pharmacol Biochem Behav* 71: 197–204.
- Glick SD, Maisonneuve IM, Raucci J, Archer S (1995). Kappa opioid inhibition of morphine and cocaine self-administration in rats. *Brain Res* 681: 147–152.
- Glick SD, Maisonneuve IM, Dickinson HA, Kitchen BA (2001). Comparative effects of dextromethorphan and dextropropanolol on morphine, methamphetamine, and nicotine self-administration in rats. *Eur J Pharmacol* 422: 87–90.
- von der Goltz C, Vengeliene V, Bilbao A, Perreau-Lenz S, Pawlak CR, Kiefer F *et al.* (2009). Cue-induced alcohol-seeking behaviour is reduced by disrupting the reconsolidation of alcohol-related memories. *Psychopharmacology (Berl)* 205: 389–397.
- Goto Y, Grace AA (2008). Limbic and cortical information processing in the nucleus accumbens. *Trends Neurosci* 31: 552–558.
- Graves SM, Napier TC (2012). SB 206553, a putative 5-HT_{2C} inverse agonist, attenuates methamphetamine-seeking in rats. *BMC Neurosci* 13: 65.
- Greco B, Carli M (2006). Reduced attention and increased impulsivity in mice lacking NPY Y2 receptors: relation to anxiolytic-like phenotype. *Behav Brain Res* 169: 325–334.
- Hadamitzky M, Koch M (2009). Effects of acute intra-cerebral administration of the 5-HT_{2A/C} receptor ligands DOI and ketanserin on impulse control in rats. *Behav Brain Res* 204: 88–92.
- Hadamitzky M, Feja M, Becker T, Koch M (2009). Effects of acute systemic administration of serotonin_{2A/C} receptor ligands in a delay-based decision-making task in rats. *Behav Pharmacol* 20: 415–423.
- Hammeress P, Joshi G, Doyle R, Georgiopoulos A, Geller D, Spencer T *et al.* (2013). Do stimulants reduce the risk for cigarette smoking in youth with attention-deficit hyperactivity disorder? A prospective, long-term, open-label study of extended-release methylphenidate. *J Pediatr* 162: 22–27 e22.
- Haney M, Spealman R (2008). Controversies in translational research: drug self-administration. *Psychopharmacology (Berl)* 199: 403–419.
- Harris GC, Hedaya MA, Pan WJ, Kalivas P (1996). beta-adrenergic antagonism alters the behavioral and neurochemical responses to cocaine. *Neuropsychopharmacology* 14: 195–204.
- Harrison AA, Everitt BJ, Robbins TW (1997). Central 5-HT depletion enhances impulsive responding without affecting the accuracy of attentional performance: interactions with dopaminergic mechanisms. *Psychopharmacology (Berl)* 133: 329–342.
- Harrison AA, Everitt BJ, Robbins TW (1999). Central serotonin depletion impairs both the acquisition and performance of a symmetrically reinforced go/no-go conditional visual discrimination. *Behav Brain Res* 100: 99–112.
- Harvey RC, Sen S, Deaciuc A, Dwoskin LP, Kantak KM (2011). Methylphenidate treatment in adolescent rats with an attention deficit/hyperactivity disorder phenotype: cocaine addiction vulnerability and dopamine transporter function. *Neuropsychopharmacology* 36: 837–847.
- Hayes DJ, Greenshaw AJ (2011). 5-HT receptors and reward-related behaviour: a review. *Neurosci Biobehav Rev* 35: 1419–1449.
- Hayes DJ, Jupp B, Sawiak SJ, Merlo E, Caprioli D, Dalley JW (2014). Brain gamma-aminobutyric acid: a neglected role in impulsivity. *Eur J Neurosci* 39: 1921–1932.
- Heidbreder CA, Andreoli M, Marcon C, Hutcheson DM, Gardner EL, Ashby CR Jr (2007). Evidence for the role of dopamine D3 receptors in oral operant alcohol self-administration and reinstatement of alcohol-seeking behavior in mice. *Addict Biol* 12: 35–50.
- Helms CM, Reeves JM, Mitchell SH (2006). Impact of strain and D-amphetamine on impulsivity (delay discounting) in inbred mice. *Psychopharmacology (Berl)* 188: 144–151.
- Higgins GA, Enderlin M, Haman M, Fletcher PJ (2003). The 5-HT_{2A} receptor antagonist M100,907 attenuates motor and ‘impulsive-type’ behaviours produced by NMDA receptor antagonism. *Psychopharmacology (Berl)* 170: 309–319.
- Higgins GA, Ballard TM, Enderlin M, Haman M, Kemp JA (2005). Evidence for improved performance in cognitive tasks following selective NR2B NMDA receptor antagonist pre-treatment in the rat. *Psychopharmacology (Berl)* 179: 85–98.
- Higgins GA, Silenies LB, Rossmann A, Rizo Z, Noble K, Soko AD *et al.* (2012). The 5-HT_{2C} receptor agonist lorcaserin reduces nicotine self-administration, discrimination, and reinstatement: relationship to feeding behavior and impulse control. *Neuropsychopharmacology* 37: 1177–1191.
- Higley AE, Kiefer SW, Li X, Gaal J, Xi ZX, Gardner EL (2011). Dopamine D(3) receptor antagonist SB-277011A inhibits

- methamphetamine self-administration and methamphetamine-induced reinstatement of drug-seeking in rats. *Eur J Pharmacol* 659: 187–192.
- Hiranita T, Soto PL, Newman AH, Katz JL (2009). Assessment of reinforcing effects of bupropion analogs and their effects on cocaine self-administration in rats: comparisons with monoamine uptake inhibitors. *J Pharmacol Exp Ther* 329: 677–686.
- Hiranita T, Soto PL, Kohut SJ, Kopajtic T, Cao J, Newman AH *et al.* (2011). Decreases in cocaine self-administration with dual inhibition of the dopamine transporter and sigma receptors. *J Pharmacol Exp Ther* 339: 662–677.
- Hobson BD, O'Neill CE, Levis SC, Monteggia LM, Neve RL, Self DW *et al.* (2013). Adenosine A1 and dopamine D1 receptor regulation of AMPA receptor phosphorylation and cocaine-seeking behavior. *Neuropsychopharmacology* 38: 1974–1983.
- Hubner CB, Moreton JE (1991). Effects of selective D1 and D2 dopamine antagonists on cocaine self-administration in the rat. *Psychopharmacology (Berl)* 105: 151–156.
- Humpston CS, Wood CM, Robinson ES (2013). Investigating the roles of different monoamine transmitters and impulse control using the 5-choice serial reaction time task. *J Psychopharmacol* 27: 213–221.
- Hyytia P, Backstrom P, Liljequist S (1999). Site-specific NMDA receptor antagonists produce differential effects on cocaine self-administration in rats. *Eur J Pharmacol* 378: 9–16.
- Isles AR, Humby T, Wilkinson LS (2003). Measuring impulsivity in mice using a novel operant delayed reinforcement task: effects of behavioural manipulations and d-amphetamine. *Psychopharmacology (Berl)* 170: 376–382.
- Ismayilova N, Shoaib M (2010). Alteration of intravenous nicotine self-administration by opioid receptor agonist and antagonists in rats. *Psychopharmacology (Berl)* 210: 211–220.
- Izzo E, Orsini C, Koob GF, Pulvirenti L (2001). A dopamine partial agonist and antagonist block amphetamine self-administration in a progressive ratio schedule. *Pharmacol Biochem Behav* 68: 701–708.
- Jayarajan P, Nirogi R, Shinde A (2013). Effect of olanzapine on scopolamine induced deficits in differential reinforcement of low rate 72s (DRL-72s) schedule in rats: involvement of the serotonergic receptors in restoring the deficits. *Eur J Pharmacol* 720: 344–354.
- Jentsch JD, Taylor JR (1999). Impulsivity resulting from frontostriatal dysfunction in drug abuse: implications for the control of behavior by reward-related stimuli. *Psychopharmacology (Berl)* 146: 373–390.
- Jupp B, Dalley JW (2014). Behavioral endophenotypes of drug addiction: etiological insights from neuroimaging studies. *Neuropharmacology* 76 (Pt B): 487–497.
- Jupp B, Lawrence AJ (2010). New horizons for therapeutics in drug and alcohol abuse. *Pharmacol Ther* 125: 138–168.
- Jupp B, Caprioli D, Dalley JW (2013a). Highly impulsive rats: modelling an endophenotype to determine the neurobiological, genetic and environmental mechanisms of addiction. *Dis Model Mech* 6: 302–311.
- Jupp B, Caprioli D, Saigal N, Reverte I, Shrestha S, Cumming P *et al.* (2013b). Dopaminergic and GABA-ergic markers of impulsivity in rats: evidence for anatomical localisation in ventral striatum and prefrontal cortex. *Eur J Neurosci* 37: 1519–1528.
- Kalivas PW, Volkow ND (2011). New medications for drug addiction hiding in glutamatergic neuroplasticity. *Mol Psychiatry* 16: 974–986.
- Kameda G, Dadmarz M, Vogel WH (2000). Influence of various drugs on the voluntary intake of nicotine by rats. *Neuropsychobiology* 41: 205–209.
- Kapur S, Remington G (1996). Serotonin-dopamine interaction and its relevance to schizophrenia. *Am J Psychiatry* 153: 466–476.
- Kayir H, Semenova S, Markou A (2014). Baseline impulsive choice predicts the effects of nicotine and nicotine withdrawal on impulsivity in rats. *Prog Neuropsychopharmacol Biol Psychiatry* 48: 6–13.
- Kertzman S, Grinspan H, Birger M, Kotler M (2006). Computerized neuropsychological examination of impulsiveness: a selective review. *Isr J Psychiatry Relat Sci* 43: 74–80.
- Khantzian EJ (1985). The self-medication hypothesis of addictive disorders: focus on heroin and cocaine dependence. *Am J Psychiatry* 142: 1259–1264.
- Khroyan TV, Barrett-Larimore RL, Rowlett JK, Spealman RD (2000). Dopamine D1- and D2-like receptor mechanisms in relapse to cocaine-seeking behavior: effects of selective antagonists and agonists. *J Pharmacol Exp Ther* 294: 680–687.
- Kirby LG, Zeeb FD, Winstanley CA (2011). Contributions of serotonin in addiction vulnerability. *Neuropharmacology* 61: 421–432.
- Kiyatkin EA, Brown PL (2003). Naloxone depresses cocaine self-administration and delays its initiation on the following day. *Neuroreport* 14: 251–255.
- Koffarnus MN, Newman AH, Grundt P, Rice KC, Woods JH (2011). Effects of selective dopaminergic compounds on a delay-discounting task. *Behav Pharmacol* 22: 300–311.
- Kolokotroni KZ, Rodgers RJ, Harrison AA (2011). Acute nicotine increases both impulsive choice and behavioural disinhibition in rats. *Psychopharmacology (Berl)* 217: 455–473.
- Kolokotroni KZ, Rodgers RJ, Harrison AA (2014). Trait differences in response to chronic nicotine and nicotine withdrawal in rats. *Psychopharmacology (Berl)* 231: 567–580.
- Koob GF (2008). A role for brain stress systems in addiction. *Neuron* 59: 11–34.
- Koob GF, Le HT, Creese I (1987). The D1 dopamine receptor antagonist SCH 23390 increases cocaine self-administration in the rat. *Neurosci Lett* 79: 315–320.
- Koskinen T, Sirvio J (2001). Studies on the involvement of the dopaminergic system in the 5-HT2 agonist (DOI)-induced premature responding in a five-choice serial reaction time task. *Brain Res Bull* 54: 65–75.
- Koskinen T, Ruotsalainen S, Puumala T, Lappalainen R, Koivisto E, Mannisto PT *et al.* (2000). Activation of 5-HT2A receptors impairs response control of rats in a five-choice serial reaction time task. *Neuropharmacology* 39: 471–481.
- Koskinen T, Haapalinna A, Sirvio J (2003). Alpha-adrenoceptor-mediated modulation of 5-HT2 receptor agonist induced impulsive responding in a 5-choice serial reaction time task. *Pharmacol Toxicol* 92: 214–225.
- Kranz GS, Kasper S, Lanzenberger R (2010). Reward and the serotonergic system. *Neuroscience* 166: 1023–1035.
- Krebs CA, Anderson KG (2012). Preference reversals and effects of D-amphetamine on delay discounting in rats. *Behav Pharmacol* 23: 228–240.
- Kuzmin A, Jerlhag E, Liljequist S, Engel J (2009). Effects of subunit selective nACh receptors on operant ethanol self-administration and relapse-like ethanol-drinking behavior. *Psychopharmacology (Berl)* 203: 99–108.

- Lacosta S, Roberts DC (1993). MDL 72222, ketanserin, and methysergide pretreatments fail to alter breaking points on a progressive ratio schedule reinforced by intravenous cocaine. *Pharmacol Biochem Behav* 44: 161–165.
- Lamb RJ, Jarbe TU (2001). Effects of fluvoxamine on ethanol-reinforced behavior in the rat. *J Pharmacol Exp Ther* 297: 1001–1009.
- Lane JD, Pickering CL, Hooper ML, Fagan K, Tyers MB, Emmett-Oglesby MW (1992). Failure of ondansetron to block the discriminative or reinforcing stimulus effects of cocaine in the rat. *Drug Alcohol Depend* 30: 151–162.
- Lane RF, Blaha CD, Rivet JM (1988). Selective inhibition of mesolimbic dopamine release following chronic administration of clozapine: involvement of alpha 1-noradrenergic receptors demonstrated by *in vivo* voltammetry. *Brain Res* 460: 398–401.
- Lasseter HC, Xie X, Arguello AA, Wells AM, Hodges MA, Fuchs RA (2014). Contribution of a mesocorticolimbic subcircuit to drug context-induced reinstatement of cocaine-seeking behavior in rats. *Neuropsychopharmacology* 39: 660–669.
- Le AD, Corrigan WA, Harding JW, Juzytsch W, Li TK (2000). Involvement of nicotinic receptors in alcohol self-administration. *Alcohol Clin Exp Res* 24: 155–163.
- Le AD, Funk D, Harding S, Juzytsch W, Li Z, Fletcher PJ (2008). Intra-median raphe nucleus (MRN) infusions of muscimol, a GABA-A receptor agonist, reinstate alcohol seeking in rats: role of impulsivity and reward. *Psychopharmacology (Berl)* 195: 605–615.
- Le AD, Harding S, Juzytsch W, Funk D, Shaham Y (2005). Role of alpha-2 adrenoceptors in stress-induced reinstatement of alcohol seeking and alcohol self-administration in rats. *Psychopharmacology (Berl)* 179: 366–373.
- Le AD, Funk D, Harding S, Juzytsch W, Fletcher PJ, Shaham Y (2006). Effects of dexfenfluramine and 5-HT₃ receptor antagonists on stress-induced reinstatement of alcohol seeking in rats. *Psychopharmacology (Berl)* 186: 82–92.
- Le AD, Funk D, Juzytsch W, Coen K, Navarre BM, Cifani C *et al.* (2011). Effect of prazosin and guanfacine on stress-induced reinstatement of alcohol and food seeking in rats. *Psychopharmacology (Berl)* 218: 89–99.
- Le Merrer J, Becker JA, Befort K, Kieffer BL (2009). Reward processing by the opioid system in the brain. *Physiol Rev* 89: 1379–1412.
- Leri F, Flores J, Rodaros D, Stewart J (2002). Blockade of stress-induced but not cocaine-induced reinstatement by infusion of noradrenergic antagonists into the bed nucleus of the stria terminalis or the central nucleus of the amygdala. *J Neurosci* 22: 5713–5718.
- Levin ED, Mead T, Rezvani AH, Rose JE, Gallivan C, Gross R (2000). The nicotinic antagonist mecamylamine preferentially inhibits cocaine vs. food self-administration in rats. *Physiol Behav* 71: 565–570.
- Levin ED, Slade S, Johnson M, Petro A, Horton K, Williams P *et al.* (2008). Ketanserin, a 5-HT₂ receptor antagonist, decreases nicotine self-administration in rats. *Eur J Pharmacol* 600: 93–97.
- Levin FR, Mariani JJ, Secora A, Brooks D, Cheng WY, Bisaga A *et al.* (2009). Atomoxetine treatment for cocaine abuse and adult attention-deficit hyperactivity disorder (ADHD): a preliminary open trial. *J Dual Diagn* 5: 41–56.
- Levy AD, Murphy JM, McBride WJ, Lumeng L, Li TK (1991). Microinjection of sulpiride into the nucleus accumbens increases ethanol drinking in alcohol-preferring (P) rats. *Alcohol Alcohol Suppl* 1: 417–420.
- Li S, Zou S, Coen K, Funk D, Shram MJ, Le AD (2014). Sex differences in yohimbine-induced increases in the reinforcing efficacy of nicotine in adolescent rats. *Addict Biol* 19: 156–164.
- Li X, Li J, Gardner EL, Xi ZX (2010). Activation of mGluR7s inhibits cocaine-induced reinstatement of drug-seeking behavior by a nucleus accumbens glutamate-mGluR2/3 mechanism in rats. *J Neurochem* 114: 1368–1380.
- Lintunen M, Hyytia P, Sallmen T, Karlstedt K, Tuomisto L, Leurs R *et al.* (2001). Increased brain histamine in an alcohol-preferring rat line and modulation of ethanol consumption by H(3) receptor mechanisms. *FASEB J* 15: 1074–1076.
- Liu X (2014). Effects of blockade of alpha4beta2 and alpha7 nicotinic acetylcholine receptors on cue-induced reinstatement of nicotine-seeking behaviour in rats. *Int J Neuropsychopharmacol* 17: 105–116.
- Liu X, Caggiula AR, Yee SK, Nobuta H, Sved AF, Pechnick RN *et al.* (2007). Mecamylamine attenuates cue-induced reinstatement of nicotine-seeking behavior in rats. *Neuropsychopharmacology* 32: 710–718.
- Liu X, Caggiula AR, Palmatier MI, Donny EC, Sved AF (2008). Cue-induced reinstatement of nicotine-seeking behavior in rats: effect of bupropion, persistence over repeated tests, and its dependence on training dose. *Psychopharmacology (Berl)* 196: 365–375.
- Liu X, Jernigen C, Gharib M, Booth S, Caggiula AR, Sved AF (2010). Effects of dopamine antagonists on drug cue-induced reinstatement of nicotine-seeking behavior in rats. *Behav Pharmacol* 21: 153–160.
- Liu YP, Lin YL, Chuang CH, Kao YC, Chang ST, Tung CS (2009). Alpha adrenergic modulation on effects of norepinephrine transporter inhibitor reboxetine in five-choice serial reaction time task. *J Biomed Sci* 16: 72.
- Logrip ML, Koob GF, Zorrilla EP (2011). Role of corticotropin-releasing factor in drug addiction: potential for pharmacological intervention. *CNS Drugs* 25: 271–287.
- Loiseau F, Le Bihan C, Hamon M, Thiebot MH (2005). Antidepressant-like effects of agomelatine, melatonin and the NK1 receptor antagonist GR205171 in impulsive-related behaviour in rats. *Psychopharmacology (Berl)* 182: 24–32.
- Lombardo LE, Bearden CE, Barrett J, Brumbaugh MS, Pittman B, Frangou S *et al.* (2012). Trait impulsivity as an endophenotype for bipolar I disorder. *Bipolar Disord* 14: 565–570.
- Loos M, Pattij T, Janssen MC, Crounotte DS, Schoffelmeer AN, Smit AB *et al.* (2010). Dopamine receptor D1/D5 gene expression in the medial prefrontal cortex predicts impulsive choice in rats. *Cereb Cortex* 20: 1064–1070.
- Lyness WH, Friedle NM, Moore KE (1980). Increased self-administration of d-amphetamine after destruction of 5-hydroxytryptaminergic neurons. *Pharmacol Biochem Behav* 12: 937–941.
- Maldonado R, Robledo P, Chover AJ, Caine SB, Koob GF (1993). D1 dopamine receptors in the nucleus accumbens modulate cocaine self-administration in the rat. *Pharmacol Biochem Behav* 45: 239–242.
- Maldonado R, Valverde O, Berrendero F (2006). Involvement of the endocannabinoid system in drug addiction. *Trends Neurosci* 29: 225–232.
- Marek GJ (2012). Activation of adenosine(1) receptors induces antidepressant-like, anti-impulsive effects on differential reinforcement of low-rate 72-s behavior in rats. *J Pharmacol Exp Ther* 341: 564–570.

- Marinelli PW, Funk D, Juzytsch W, Harding S, Rice KC, Shaham Y *et al.* (2007). The CRF1 receptor antagonist antalarmin attenuates yohimbine-induced increases in operant alcohol self-administration and reinstatement of alcohol seeking in rats. *Psychopharmacology (Berl)* 195: 345–355.
- Marinelli PW, Funk D, Harding S, Li Z, Juzytsch W, Le AD (2009). Roles of opioid receptor subtypes in mediating alcohol-seeking induced by discrete cues and context. *Eur J Neurosci* 30: 671–678.
- Martelle SE, Porrino LJ, Nader MA (2013). Effects of chronic methylphenidate in adolescence on later methylphenidate self-administration in rhesus monkeys. *Behav Pharmacol* 24: 478–481.
- Marusich JA, Bardo MT (2009). Differences in impulsivity on a delay-discounting task predict self-administration of a low unit dose of methylphenidate in rats. *Behav Pharmacol* 20: 447–454.
- Maurel S, De Vry J, Schreiber R (1999). 5-HT receptor ligands differentially affect operant oral self-administration of ethanol in the rat. *Eur J Pharmacol* 370: 217–223.
- McKinzie DL, McBride WJ, Murphy JM, Lumeng L, Li TK (2000). Effects of MDL 72222, a serotonin₃ antagonist, on operant responding for ethanol by alcohol-preferring P rats. *Alcohol Clin Exp Res* 24: 1500–1504.
- Mendez IA, Simon NW, Hart N, Mitchell MR, Nation JR, Wellman PJ *et al.* (2010). Self-administered cocaine causes long-lasting increases in impulsive choice in a delay discounting task. *Behav Neurosci* 124: 470–477.
- Mendez IA, Gilbert RJ, Bizon JL, Setlow B (2012). Effects of acute administration of nicotinic and muscarinic cholinergic agonists and antagonists on performance in different cost–benefit decision making tasks in rats. *Psychopharmacology (Berl)* 224: 489–499.
- Milstein JA, Lehmann O, Theobald DE, Dalley JW, Robbins TW (2007). Selective depletion of cortical noradrenaline by anti-dopamine beta-hydroxylase-saporin impairs attentional function and enhances the effects of guanfacine in the rat. *Psychopharmacology (Berl)* 190: 51–63.
- Milstein JA, Dalley JW, Robbins TW (2010). Methylphenidate-induced impulsivity: pharmacological antagonism by beta-adrenoreceptor blockade. *J Psychopharmacol* 24: 309–321.
- Mirza NR, Stolerman IP (2000). The role of nicotinic and muscarinic acetylcholine receptors in attention. *Psychopharmacology (Berl)* 148: 243–250.
- Miyazaki K, Miyazaki KW, Doya K (2012). The role of serotonin in the regulation of patience and impulsivity. *Mol Neurobiol* 45: 213–224.
- Mobini S, Chiang TJ, Ho MY, Bradshaw CM, Szabadi E (2000). Effects of central 5-hydroxytryptamine depletion on sensitivity to delayed and probabilistic reinforcement. *Psychopharmacology (Berl)* 152: 390–397.
- Moeller FG, Barratt ES, Dougherty DM, Schmitz JM, Swann AC (2001). Psychiatric aspects of impulsivity. *Am J Psychiatry* 158: 1783–1793.
- Molander AC, Mar A, Norbury A, Steventon S, Moreno M, Caprioli D *et al.* (2011). High impulsivity predicting vulnerability to cocaine addiction in rats: some relationship with novelty preference but not novelty reactivity, anxiety or stress. *Psychopharmacology (Berl)* 215: 721–731.
- Monterosso J, Ainslie G (1999). Beyond discounting: possible experimental models of impulse control. *Psychopharmacology (Berl)* 146: 339–347.
- Moreno-Lopez L, Catena A, Fernandez-Serrano MJ, Delgado-Rico E, Stamatakis EA, Perez-Garcia M *et al.* (2012). Trait impulsivity and prefrontal gray matter reductions in cocaine dependent individuals. *Drug Alcohol Depend* 125: 208–214.
- Muller SE, Weijers HG, Boning J, Wiesbeck GA (2008). Personality traits predict treatment outcome in alcohol-dependent patients. *Neuropsychobiology* 57: 159–164.
- Munzar P, Tanda G, Justinova Z, Goldberg SR (2004). Histamine h₃ receptor antagonists potentiate methamphetamine self-administration and methamphetamine-induced accumbal dopamine release. *Neuropsychopharmacology* 29: 705–717.
- Murnane KS, Winschel J, Schmidt KT, Stewart LM, Rose SJ, Cheng K *et al.* (2013). Serotonin 2A receptors differentially contribute to abuse-related effects of cocaine and cocaine-induced nigrostriatal and mesolimbic dopamine overflow in nonhuman primates. *J Neurosci* 33: 13367–13374.
- Murphy ER, Fernando AB, Urcelay GP, Robinson ES, Mar AC, Theobald DE *et al.* (2012). Impulsive behaviour induced by both NMDA receptor antagonism and GABAA receptor activation in rat ventromedial prefrontal cortex. *Psychopharmacology (Berl)* 219: 401–410.
- Navarra R, Comery TA, Graf R, Rosenzweig-Lipson S, Day M (2008a). The 5-HT_{2C} receptor agonist WAY-163909 decreases impulsivity in the 5-choice serial reaction time test. *Behav Brain Res* 188: 412–415.
- Navarra R, Graf R, Huang Y, Logue S, Comery T, Hughes Z *et al.* (2008b). Effects of atomoxetine and methylphenidate on attention and impulsivity in the 5-choice serial reaction time test. *Prog Neuropsychopharmacol Biol Psychiatry* 32: 34–41.
- Nemeth CL, Paine TA, Rittiner JE, Beguin C, Carroll FI, Roth BL *et al.* (2010). Role of kappa-opioid receptors in the effects of salvinorin A and ketamine on attention in rats. *Psychopharmacology (Berl)* 210: 263–274.
- Nestler EJ (2005). The neurobiology of cocaine addiction. *Sci Pract Perspect* 3: 4–10.
- Nic Dhonnchadha BA, Fox RG, Stutz SJ, Rice KC, Cunningham KA (2009). Blockade of the serotonin 5-HT_{2A} receptor suppresses cue-evoked reinstatement of cocaine-seeking behavior in a rat self-administration model. *Behav Neurosci* 123: 382–396.
- Nigg JT, Wong MM, Martel MM, Jester JM, Puttler LJ, Glass JM *et al.* (2006). Poor response inhibition as a predictor of problem drinking and illicit drug use in adolescents at risk for alcoholism and other substance use disorders. *J Am Acad Child Adolesc Psychiatry* 45: 468–475.
- Nuutinen S, Lintunen M, Vanhanen J, Ojala T, Rozov S, Panula P (2011). Evidence for the role of histamine H₃ receptor in alcohol consumption and alcohol reward in mice. *Neuropsychopharmacology* 36: 2030–2040.
- Oberlin BG, Grahame NJ (2009). High-alcohol preferring mice are more impulsive than low-alcohol preferring mice as measured in the delay discounting task. *Alcohol Clin Exp Res* 33: 1294–1303.
- Oberlin BG, Bristow RE, Heighton ME, Grahame NJ (2010). Pharmacologic dissociation between impulsivity and alcohol drinking in high alcohol preferring mice. *Alcohol Clin Exp Res* 34: 1363–1375.
- O'Brien CP (2008). Review. Evidence-based treatments of addiction. *Philos Trans R Soc Lond B Biol Sci* 363: 3277–3286.
- Ohmura Y, Yamaguchi T, Futami Y, Togashi H, Izumi T, Matsumoto M *et al.* (2009). Corticotropin releasing factor enhances attentional

function as assessed by the five-choice serial reaction time task in rats. *Behav Brain Res* 198: 429–433.

Oliver YP, Ripley TL, Stephens DN (2009). Ethanol effects on impulsivity in two mouse strains: similarities to diazepam and ketamine. *Psychopharmacology (Berl)* 204: 679–692.

Olmstead MC, Hellems KG, Paine TA (2006). Alcohol-induced impulsivity in rats: an effect of cue salience? *Psychopharmacology (Berl)* 184: 221–228.

Olsen CM, Duvauchelle CL (2006). Prefrontal cortex D1 modulation of the reinforcing properties of cocaine. *Brain Res* 1075: 229–235.

Paine TA, Dringenberg HC, Olmstead MC (2003). Effects of chronic cocaine on impulsivity: relation to cortical serotonin mechanisms. *Behav Brain Res* 147: 135–147.

Paine TA, Tomasiewicz HC, Zhang K, Carlezon WA Jr (2007). Sensitivity of the five-choice serial reaction time task to the effects of various psychotropic drugs in Sprague-Dawley rats. *Biol Psychiatry* 62: 687–693.

Palmatier MI, Liu X, Donny EC, Caggiula AR, Sved AF (2008). Metabotropic glutamate 5 receptor (mGluR5) antagonists decrease nicotine seeking, but do not affect the reinforcement enhancing effects of nicotine. *Neuropsychopharmacology* 33: 2139–2147.

Pardey MC, Kumar NN, Goodchild AK, Cornish JL (2013). Catecholamine receptors differentially mediate impulsive choice in the medial prefrontal and orbitofrontal cortex. *J Psychopharmacol* 27: 203–212.

Parsons LH, Weiss F, Koob GF (1998). Serotonin1B receptor stimulation enhances cocaine reinforcement. *J Neurosci* 18: 10078–10089.

Passetti F, Dalley JW, Robbins TW (2003). Double dissociation of serotonergic and dopaminergic mechanisms on attentional performance using a rodent five-choice reaction time task. *Psychopharmacology (Berl)* 165: 136–145.

Paterson NE, Markou A (2005). The metabotropic glutamate receptor 5 antagonist MPEP decreased break points for nicotine, cocaine and food in rats. *Psychopharmacology (Berl)* 179: 255–261.

Paterson NE, Wetzler C, Hackett A, Hanania T (2012). Impulsive action and impulsive choice are mediated by distinct neuropharmacological substrates in rat. *Int J Neuropsychopharmacol* 15: 1473–1487.

Pattij T, Vanderschuren LJ (2008). The neuropharmacology of impulsive behaviour. *Trends Pharmacol Sci* 29: 192–199.

Pattij T, Janssen MC, Vanderschuren LJ, Schoffeleer AN, van Gaalen MM (2007a). Involvement of dopamine D1 and D2 receptors in the nucleus accumbens core and shell in inhibitory response control. *Psychopharmacology (Berl)* 191: 587–598.

Pattij T, Janssen MC, Schepers I, Gonzalez-Cuevas G, de Vries TJ, Schoffeleer AN (2007b). Effects of the cannabinoid CB1 receptor antagonist rimonabant on distinct measures of impulsive behavior in rats. *Psychopharmacology (Berl)* 193: 85–96.

Pattij T, Schetters D, Janssen MC, Wiskerke J, Schoffeleer AN (2009). Acute effects of morphine on distinct forms of impulsive behavior in rats. *Psychopharmacology (Berl)* 205: 489–502.

Pattij T, Schetters D, Schoffeleer AN, van Gaalen MM (2012). On the improvement of inhibitory response control and visuospatial attention by indirect and direct adrenoceptor agonists. *Psychopharmacology (Berl)* 219: 327–340.

Pawson AJ, Sharman JL, Benson HE, Faccenda E, Alexander SP, Buneman OP *et al.*; NC-IUPHAR (2014). The IUPHAR/BPS Guide to

PHARMACOLOGY: an expert-driven knowledgebase of drug targets and their ligands. *Nucl. Acids Res* 42 (Database Issue): D1098–106.

Pelloux Y, Dilleen R, Economidou D, Theobald D, Everitt BJ (2012). Reduced forebrain serotonin transmission is causally involved in the development of compulsive cocaine seeking in rats. *Neuropsychopharmacology* 37: 2505–2514.

Peltier R, Schenk S (1993). Effects of serotonergic manipulations on cocaine self-administration in rats. *Psychopharmacology (Berl)* 110: 390–394.

Pentkowski NS, Duke FD, Weber SM, Pockros LA, Teer AP, Hamilton EC *et al.* (2010). Stimulation of medial prefrontal cortex serotonin 2C (5-HT_{2C}) receptors attenuates cocaine-seeking behavior. *Neuropsychopharmacology* 35: 2037–2048.

Perry JC, Korner AC (2011). Impulsive phenomena, the impulsive character (der Triebhafte Charakter) and DSM personality disorders. *J Pers Disord* 25: 586–606.

Perry JL, Stairs DJ, Bardo MT (2008). Impulsive choice and environmental enrichment: effects of d-amphetamine and methylphenidate. *Behav Brain Res* 193: 48–54.

Petry NM (2001). Delay discounting of money and alcohol in actively using alcoholics, currently abstinent alcoholics, and controls. *Psychopharmacology (Berl)* 154: 243–250.

Pezze MA, Dalley JW, Robbins TW (2007). Differential roles of dopamine D1 and D2 receptors in the nucleus accumbens in attentional performance on the five-choice serial reaction time task. *Neuropsychopharmacology* 32: 273–283.

Pfeffer AO, Samson HH (1985). Oral ethanol reinforcement in the rat: effects of acute amphetamine. *Alcohol* 2: 693–697.

Phillips GD, Howes SR, Whitelaw RB, Robbins TW, Everitt BJ (1994). Isolation rearing impairs the reinforcing efficacy of intravenous cocaine or intra-accumbens d-amphetamine: impaired response to intra-accumbens D1 and D2/D3 dopamine receptor antagonists. *Psychopharmacology (Berl)* 115: 419–429.

Pierce RC, O'Brien CP, Kenny PJ, Vanderschuren LJ (2012). Rational development of addiction pharmacotherapies: successes, failures, and prospects. *Cold Spring Harb Perspect Med* 2: a012880.

Placenza FM, Fletcher PJ, Vaccarino FJ, Erb S (2006). Effects of central neurokinin-1 receptor antagonism on cocaine- and opiate-induced locomotor activity and self-administration behaviour in rats. *Pharmacol Biochem Behav* 84: 94–101.

Poulos CX, Le AD, Parker JL (1995). Impulsivity predicts individual susceptibility to high levels of alcohol self-administration. *Behav Pharmacol* 6: 810–814.

Przegalinski E, Golda A, Filip M (2008). Effects of serotonin (5-HT) (1B) receptor ligands on cocaine-seeking behavior in rats. *Pharmacol Rep* 60: 798–810.

Radwanska K, Kaczmarek L (2012). Characterization of an alcohol addiction-prone phenotype in mice. *Addict Biol* 17: 601–612.

Rasmussen DD, Alexander LL, Raskind MA, Froehlich JC (2009). The alpha1-adrenergic receptor antagonist, prazosin, reduces alcohol drinking in alcohol-preferring (P) rats. *Alcohol Clin Exp Res* 33: 264–272.

Rasmussen T, Sauerberg P, Nielsen EB, Swedberg MD, Thomsen C, Sheardown MJ *et al.* (2000). Muscarinic receptor agonists decrease cocaine self-administration rates in drug-naïve mice. *Eur J Pharmacol* 402: 241–246.

Rauhut AS, Mullins SN, Dwoskin LP, Bardo MT (2002). Reboxetine: attenuation of intravenous nicotine self-administration in rats. *J Pharmacol Exp Ther* 303: 664–672.

- Redila VA, Chavkin C (2008). Stress-induced reinstatement of cocaine seeking is mediated by the kappa opioid system. *Psychopharmacology (Berl)* 200: 59–70.
- van Ree JM, Gerrits MA, Vanderschuren LJ (1999). Opioids, reward and addiction: an encounter of biology, psychology, and medicine. *Pharmacol Rev* 51: 341–396.
- Reichel CM, Murray JE, Grant KM, Bevins RA (2009). Bupropion attenuates methamphetamine self-administration in adult male rats. *Drug Alcohol Depend* 100: 54–62.
- Reynolds B, de Wit H, Richards J (2002). Delay of gratification and delay discounting in rats. *Behav Processes* 59: 157.
- Rezvani AH, Overstreet DH, Janowsky DS (1991). Drug-induced reductions in ethanol intake in alcohol preferring and Fawn-Hooded rats. *Alcohol Alcohol Suppl* 1: 433–437.
- Richards JB, Zhang L, Mitchell SH, de Wit H (1999). Delay or probability discounting in a model of impulsive behavior: effect of alcohol. *J Exp Anal Behav* 71: 121–143.
- Richardson NR, Smith AM, Roberts DC (1994). A single injection of either flupenthixol decanoate or haloperidol decanoate produces long-term changes in cocaine self-administration in rats. *Drug Alcohol Depend* 36: 23–25.
- Rimondini R, Thorsell A, Heilig M (2005). Suppression of ethanol self-administration by the neuropeptide Y (NPY) Y2 receptor antagonist BIIIE0246: evidence for sensitization in rats with a history of dependence. *Neurosci Lett* 375: 129–133.
- Robbins TW (2002). The 5-choice serial reaction time task: behavioural pharmacology and functional neurochemistry. *Psychopharmacology (Berl)* 163: 362–380.
- Robbins TW, Gillan CM, Smith DG, de Wit S, Ersche KD (2012). Neurocognitive endophenotypes of impulsivity and compulsivity: towards dimensional psychiatry. *Trends Cogn Sci* 16: 81–91.
- Roberts DC, Loh EA, Baker GB, Vickers G (1994). Lesions of central serotonin systems affect responding on a progressive ratio schedule reinforced either by intravenous cocaine or by food. *Pharmacol Biochem Behav* 49: 177–182.
- Robinson ES, Dalley JW, Theobald DE, Glennon JC, Pezze MA, Murphy ER *et al.* (2008a). Opposing roles for 5-HT_{2A} and 5-HT_{2C} receptors in the nucleus accumbens on inhibitory response control in the 5-choice serial reaction time task. *Neuropsychopharmacology* 33: 2398–2406.
- Robinson ES, Eagle DM, Mar AC, Bari A, Banerjee G, Jiang X *et al.* (2008b). Similar effects of the selective noradrenaline reuptake inhibitor atomoxetine on three distinct forms of impulsivity in the rat. *Neuropsychopharmacology* 33: 1028–1037.
- Roesch MR, Takahashi Y, Gugs N, Bissonette GB, Schoenbaum G (2007). Previous cocaine exposure makes rats hypersensitive to both delay and reward magnitude. *J Neurosci* 27: 245–250.
- Rogers RD, Robbins TW (2001). Investigating the neurocognitive deficits associated with chronic drug misuse. *Curr Opin Neurobiol* 11: 250–257.
- Roman V, Gyertyan I, Saghy K, Kiss B, Szombathelyi Z (2013). Cariprazine (RGH-188), a D(3)-preferring dopamine D(3)/D(2) receptor partial agonist antipsychotic candidate demonstrates anti-abuse potential in rats. *Psychopharmacology (Berl)* 226: 285–293.
- Ross JT, Corrigan WA, Heidbreder CA, LeSage MG (2007). Effects of the selective dopamine D3 receptor antagonist SB-277011A on the reinforcing effects of nicotine as measured by a progressive-ratio schedule in rats. *Eur J Pharmacol* 559: 173–179.
- Rowlett JK, Platt DM, Yao WD, Spealman RD (2007). Modulation of heroin and cocaine self-administration by dopamine D1- and D2-like receptor agonists in rhesus monkeys. *J Pharmacol Exp Ther* 321: 1135–1143.
- Ruedi-Bettschen D, Rowlett JK, Spealman RD, Platt DM (2010). Attenuation of cocaine-induced reinstatement of drug seeking in squirrel monkeys: kappa opioid and serotonergic mechanisms. *Psychopharmacology (Berl)* 210: 169–177.
- Ruotsalainen S, Miettinen R, MacDonald E, Koivisto E, Sirvio J (2000). Blockade of muscarinic, rather than nicotinic, receptors impairs attention, but does not interact with serotonin depletion. *Psychopharmacology (Berl)* 148: 111–123.
- Sabino V, Cottone P, Steardo L, Schmidhammer H, Zorrilla EP (2007). 14-Methoxymetopon, a highly potent mu opioid agonist, biphasically affects ethanol intake in Sardinian alcohol-preferring rats. *Psychopharmacology (Berl)* 192: 537–546.
- Sabino V, Narayan AR, Zeric T, Steardo L, Cottone P (2013). mTOR activation is required for the anti-alcohol effect of ketamine, but not memantine, in alcohol-preferring rats. *Behav Brain Res* 247: 9–16.
- Samson HH, Chappell A (2003). Dopaminergic involvement in medial prefrontal cortex and core of the nucleus accumbens in the regulation of ethanol self-administration: a dual-site microinjection study in the rat. *Physiol Behav* 79: 581–590.
- Samson HH, Hodge CW, Tolliver GA, Haraguchi M (1993). Effect of dopamine agonists and antagonists on ethanol-reinforced behavior: the involvement of the nucleus accumbens. *Brain Res Bull* 30: 133–141.
- Samson HH, Chappell A, Slawecki C, Hodge C (1999). The effects of microinjection of d-amphetamine into the n. accumbens during the late maintenance phase of an ethanol consumption bout. *Pharmacol Biochem Behav* 63: 159–165.
- Sanchis-Segura C, Spanagel R (2006). Behavioural assessment of drug reinforcement and addictive features in rodents: an overview. *Addict Biol* 11: 2–38.
- Schank JR, Pickens CL, Rowe KE, Cheng K, Thorsell A, Rice KC *et al.* (2011). Stress-induced reinstatement of alcohol-seeking in rats is selectively suppressed by the neurokinin 1 (NK1) antagonist L822429. *Psychopharmacology (Berl)* 218: 111–119.
- Schank JR, Ryabinin AE, Giardino WJ, Ciccocioppo R, Heilig M (2012). Stress-related neuropeptides and addictive behaviors: beyond the usual suspects. *Neuron* 76: 192–208.
- Schank JR, King CE, Sun H, Cheng K, Rice KC, Heilig M *et al.* (2014). The role of the neurokinin-1 receptor in stress-induced reinstatement of alcohol and cocaine seeking. *Neuropsychopharmacology* 39: 1093–1101.
- Schenk S (2000). Effects of the serotonin 5-HT(2) antagonist, ritanserin, and the serotonin 5-HT(1A) antagonist, WAY 100635, on cocaine-seeking in rats. *Pharmacol Biochem Behav* 67: 363–369.
- Schenk S (2002). Effects of GBR 12909, WIN 35,428 and indatraline on cocaine self-administration and cocaine seeking in rats. *Psychopharmacology (Berl)* 160: 263–270.
- Schenk S, Gittings D (2003). Effects of SCH 23390 and eticlopride on cocaine-seeking produced by cocaine and WIN 35,428 in rats. *Psychopharmacology (Berl)* 168: 118–123.
- Schenk S, Partridge B (1999). Cocaine-seeking produced by experimenter-administered drug injections: dose–effect relationships in rats. *Psychopharmacology (Berl)* 147: 285–290.
- Schenk S, Partridge B, Shippenberg TS (2000). Reinstatement of extinguished drug-taking behavior in rats: effect of the kappa-opioid receptor agonist, U69593. *Psychopharmacology (Berl)* 151: 85–90.

- Schippers MC, Binnekade R, Schoffelmeer AN, Pattij T, De Vries TJ (2012). Unidirectional relationship between heroin self-administration and impulsive decision-making in rats. *Psychopharmacology (Berl)* 219: 443–452.
- Schmidt HD, Anderson SM, Pierce RC (2006). Stimulation of D1-like or D2 dopamine receptors in the shell, but not the core, of the nucleus accumbens reinstates cocaine-seeking behaviour in the rat. *Eur J Neurosci* 23: 219–228.
- Schmidt KT, Weinshenker D (2014). Adrenaline rush: the role of adrenergic receptors in stimulant-induced behaviors. *Mol Pharmacol* 85: 640–650.
- Schroeder JP, Overstreet DH, Hodge CW (2005). The mGluR5 antagonist MPEP decreases operant ethanol self-administration during maintenance and after repeated alcohol deprivations in alcohol-preferring (P) rats. *Psychopharmacology (Berl)* 179: 262–270.
- Self D (2010). Dopamine receptor subtypes in reward and relapse. In: Neve KA (ed.). *The Dopamine Receptors*. Humana Press: New York, pp. 479–523.
- Self DW, Barnhart WJ, Lehman DA, Nestler EJ (1996). Opposite modulation of cocaine-seeking behavior by D1- and D2-like dopamine receptor agonists. *Science* 271: 1586–1589.
- Semenova S, Markou A (2007). The effects of the mGluR5 antagonist MPEP and the mGluR2/3 antagonist LY341495 on rats' performance in the 5-choice serial reaction time task. *Neuropharmacology* 52: 863–872.
- Shannon HE, Eberle EL (2006). Effects of biasing the location of stimulus presentation, and the muscarinic cholinergic receptor antagonist scopolamine, on performance of a 5-choice serial reaction time attention task in rats. *Behav Pharmacol* 17: 71–85.
- Shelton KL, Balster RL (1997). Effects of gamma-aminobutyric acid agonists and N-methyl-D-aspartate antagonists on a multiple schedule of ethanol and saccharin self-administration in rats. *J Pharmacol Exp Ther* 280: 1250–1260.
- Shepard JD, Bossert JM, Liu SY, Shaham Y (2004). The anxiogenic drug yohimbine reinstates methamphetamine seeking in a rat model of drug relapse. *Biol Psychiatry* 55: 1082–1089.
- Simmons D, Self DW (2009). Role of mu- and delta-opioid receptors in the nucleus accumbens in cocaine-seeking behavior. *Neuropsychopharmacology* 34: 1946–1957.
- Simon NW, Beas BS, Montgomery KS, Haberman RP, Bizon JL, Setlow B (2013). Prefrontal cortical-striatal dopamine receptor mRNA expression predicts distinct forms of impulsivity. *Eur J Neurosci* 37: 1779–1788.
- Simon O'Brien E, Legastelois R, Houchi H, Vilpoux C, Alaux-Cantin S, Pierrefiche O *et al.* (2011). Fluoxetine, desipramine, and the dual antidepressant milnacipran reduce alcohol self-administration and/or relapse in dependent rats. *Neuropsychopharmacology* 36: 1518–1530.
- Simpson D, Plosker GL (2004). Atomoxetine: a review of its use in adults with attention deficit hyperactivity disorder. *Drugs* 64: 205–222.
- Smith RJ, Aston-Jones G (2011). $\alpha(2)$ Adrenergic and imidazoline receptor agonists prevent cue-induced cocaine seeking. *Biol Psychiatry* 70: 712–719.
- Sofuoglu M, Sewell RA (2009). Norepinephrine and stimulant addiction. *Addict Biol* 14: 119–129.
- Solanto MV (1984). Neuropharmacological basis of stimulant drug action in attention deficit disorder with hyperactivity: a review and synthesis. *Psychol Bull* 95: 387–409.
- Somkuwar SS, Jordan CJ, Kantak KM, Dwoskin LP (2013). Adolescent atomoxetine treatment in a rodent model of ADHD: effects on cocaine self-administration and dopamine transporters in frontostriatal regions. *Neuropsychopharmacology* 38: 2588–2597.
- Song R, Yang RF, Wu N, Su RB, Li J, Peng XQ *et al.* (2012). YQA14: a novel dopamine D3 receptor antagonist that inhibits cocaine self-administration in rats and mice, but not in D3 receptor-knockout mice. *Addict Biol* 17: 259–273.
- Soria G, Mendizabal V, Tourino C, Robledo P, Ledent C, Parmentier M *et al.* (2005). Lack of CB1 cannabinoid receptor impairs cocaine self-administration. *Neuropsychopharmacology* 30: 1670–1680.
- Spano MS, Fattore L, Cossu G, Deiana S, Fadda P, Fratta W (2004). CB1 receptor agonist and heroin, but not cocaine, reinstate cannabinoid-seeking behaviour in the rat. *Br J Pharmacol* 143: 343–350.
- Stanis JJ, Marquez Avila H, White MD, Gulley JM (2008a). Dissociation between long-lasting behavioral sensitization to amphetamine and impulsive choice in rats performing a delay-discounting task. *Psychopharmacology (Berl)* 199: 539–548.
- Stanis JJ, Burns RM, Sherrill LK, Gulley JM (2008b). Disparate cocaine-induced locomotion as a predictor of choice behavior in rats trained in a delay-discounting task. *Drug Alcohol Depend* 98: 54–62.
- Sukhotina IA, Dravolina OA, Novitskaya Y, Zvartau EE, Danysz W, Bessalov AY (2008). Effects of mGlu1 receptor blockade on working memory, time estimation, and impulsivity in rats. *Psychopharmacology (Berl)* 196: 211–220.
- Sulzer D (2011). How addictive drugs disrupt presynaptic dopamine neurotransmission. *Neuron* 69: 628–649.
- Sun H, Green TA, Theobald DE, Birnbaum SG, Graham DL, Zeeb FD *et al.* (2010). Yohimbine increases impulsivity through activation of cAMP response element binding in the orbitofrontal cortex. *Biol Psychiatry* 67: 649–656.
- Sun W, Rebec GV (2005). The role of prefrontal cortex D1-like and D2-like receptors in cocaine-seeking behavior in rats. *Psychopharmacology (Berl)* 177: 315–323.
- Suto N, Wise RA (2011). Satiating effects of cocaine are controlled by dopamine actions in the nucleus accumbens core. *J Neurosci* 31: 17917–17922.
- Szerman N, Peris L, Mesias B, Colis P, Rosa J, Prieto A (2005). Reboxetine for the treatment of patients with Cocaine Dependence Disorder. *Hum Psychopharmacol* 20: 189–192.
- Talpos JC, Wilkinson LS, Robbins TW (2006). A comparison of multiple 5-HT receptors in two tasks measuring impulsivity. *J Psychopharmacol* 20: 47–58.
- Tella SR (1995). Effects of monoamine reuptake inhibitors on cocaine self-administration in rats. *Pharmacol Biochem Behav* 51: 687–692.
- Tessari M, Pilla M, Andreoli M, Hutcheson DM, Heidbreder CA (2004). Antagonism at metabotropic glutamate 5 receptors inhibits nicotine- and cocaine-taking behaviours and prevents nicotine-triggered relapse to nicotine-seeking. *Eur J Pharmacol* 499: 121–133.
- Thanos PK, Michaelides M, Benveniste H, Wang GJ, Volkow ND (2007). Effects of chronic oral methylphenidate on cocaine self-administration and striatal dopamine D2 receptors in rodents. *Pharmacol Biochem Behav* 87: 426–433.
- Thiebot MH, Le Bihan C, Soubrie P, Simon P (1985). Benzodiazepines reduce the tolerance to reward delay in rats. *Psychopharmacology (Berl)* 86: 147–152.

- Thomas MJ, Kalivas PW, Shaham Y (2008). Neuroplasticity in the mesolimbic dopamine system and cocaine addiction. *Br J Pharmacol* 154: 327–342.
- Thomsen M, Conn PJ, Lindsley C, Wess J, Boon JY, Fulton BS *et al.* (2010). Attenuation of cocaine's reinforcing and discriminative stimulus effects via muscarinic M1 acetylcholine receptor stimulation. *J Pharmacol Exp Ther* 332: 959–969.
- Thomsen M, Fulton BS, Caine SB (2014). Acute and chronic effects of the M1/M4 preferring muscarinic agonist xanomeline on cocaine vs. food choice in rats. *Psychopharmacology (Berl)* 231: 469–479.
- Thorsell A, Rimondini R, Heilig M (2002). Blockade of central neuropeptide Y (NPY) Y2 receptors reduces ethanol self-administration in rats. *Neurosci Lett* 332: 1–4.
- Tomkins DM, Le AD, Sellers EM (1995). Effect of the 5-HT3 antagonist ondansetron on voluntary ethanol intake in rats and mice maintained on a limited access procedure. *Psychopharmacology (Berl)* 117: 479–485.
- Tomkins DM, Joharchi N, Tampakeras M, Martin JR, Wichmann J, Higgins GA (2002). An investigation of the role of 5-HT(2C) receptors in modifying ethanol self-administration behaviour. *Pharmacol Biochem Behav* 71: 735–744.
- Torregrossa MM, Xie M, Taylor JR (2012). Chronic corticosterone exposure during adolescence reduces impulsive action but increases impulsive choice and sensitivity to yohimbine in male Sprague-Dawley rats. *Neuropsychopharmacology* 37: 1656–1670.
- Tran-Nguyen LT, Bellew JG, Grote KA, Neisewander JL (2001). Serotonin depletion attenuates cocaine seeking but enhances sucrose seeking and the effects of cocaine priming on reinstatement of cocaine seeking in rats. *Psychopharmacology (Berl)* 157: 340–348.
- Tsutsui-Kimura I, Ohmura Y, Izumi T, Yamaguchi T, Yoshida T, Yoshioka M (2009). The effects of serotonin and/or noradrenaline reuptake inhibitors on impulsive-like action assessed by the three-choice serial reaction time task: a simple and valid model of impulsive action using rats. *Behav Pharmacol* 20: 474–483.
- Tsutsui-Kimura I, Ohmura Y, Izumi T, Yamaguchi T, Yoshida T, Yoshioka M (2010). Endogenous acetylcholine modulates impulsive action via $\alpha 4\beta 2$ nicotinic acetylcholine receptors in rats. *Eur J Pharmacol* 641: 148–153.
- Uhl GR (2006). Molecular genetics of addiction vulnerability. *NeuroRx* 3: 295–301.
- Vanderschuren LJ, Everitt BJ (2004). Drug seeking becomes compulsive after prolonged cocaine self-administration. *Science* 305: 1017–1019.
- Vengeliene V, Bachteler D, Danysz W, Spanagel R (2005). The role of the NMDA receptor in alcohol relapse: a pharmacological mapping study using the alcohol deprivation effect. *Neuropharmacology* 48: 822–829.
- Verdejo-Garcia A, Lawrence AJ, Clark L (2008). Impulsivity as a vulnerability marker for substance-use disorders: review of findings from high-risk research, problem gamblers and genetic association studies. *Neurosci Biobehav Rev* 32: 777–810.
- Volkow ND, Fowler JS, Wang GJ, Swanson JM (2004). Dopamine in drug abuse and addiction: results from imaging studies and treatment implications. *Mol Psychiatry* 9: 557–569.
- de Vries TJ, Babovic-Vuksanovic D, Elmer G, Shippenberg TS (1995). Lack of involvement of delta-opioid receptors in mediating the rewarding effects of cocaine. *Psychopharmacology (Berl)* 120: 442–448.
- Wade TR, de Wit H, Richards JB (2000). Effects of dopaminergic drugs on delayed reward as a measure of impulsive behavior in rats. *Psychopharmacology (Berl)* 150: 90–101.
- Walker BM, Kissler JL (2013). Dissociable effects of kappa-opioid receptor activation on impulsive phenotypes in wistar rats. *Neuropsychopharmacology* 38: 2278–2285.
- Walker BM, Zorrilla EP, Koob GF (2011). Systemic kappa-opioid receptor antagonism by nor-binaltorphimine reduces dependence-induced excessive alcohol self-administration in rats. *Addict Biol* 16: 116–119.
- Walsh SL, Middleton LS, Wong CJ, Nuzzo PA, Campbell CL, Rush CR *et al.* (2013). Atomoxetine does not alter cocaine use in cocaine dependent individuals: double blind randomized trial. *Drug Alcohol Depend* 130: 150–157.
- Wang J, Lanfranco MF, Gibb SL, Yowell QV, Carnicella S, Ron D (2010). Long-lasting adaptations of the NR2B-containing NMDA receptors in the dorsomedial striatum play a crucial role in alcohol consumption and relapse. *J Neurosci* 30: 10187–10198.
- Ward AS, Li DH, Luedtke RR, Emmett-Oglesby MW (1996). Variations in cocaine self-administration by inbred rat strains under a progressive-ratio schedule. *Psychopharmacology (Berl)* 127: 204–212.
- Watkins SS, Epping-Jordan MP, Koob GF, Markou A (1999). Blockade of nicotine self-administration with nicotinic antagonists in rats. *Pharmacol Biochem Behav* 62: 743–751.
- Wee S, Wang Z, Woolverton WL, Pulvirenti L, Koob GF (2007). Effect of aripiprazole, a partial dopamine D2 receptor agonist, on increased rate of methamphetamine self-administration in rats with prolonged session duration. *Neuropsychopharmacology* 32: 2238–2247.
- Wee S, Orio L, Ghirmai S, Cashman JR, Koob GF (2009). Inhibition of kappa opioid receptors attenuated increased cocaine intake in rats with extended access to cocaine. *Psychopharmacology (Berl)* 205: 565–575.
- Weinshenker D, Schroeder JP (2007). There and back again: a tale of norepinephrine and drug addiction. *Neuropsychopharmacology* 32: 1433–1451.
- Weiss F, Martin-Fardon R, Ciccocioppo R, Kerr TM, Smith DL, Ben-Shahar O (2001). Enduring resistance to extinction of cocaine-seeking behavior induced by drug-related cues. *Neuropsychopharmacology* 25: 361–372.
- Weissenborn R, Deroche V, Koob GF, Weiss F (1996). Effects of dopamine agonists and antagonists on cocaine-induced operant responding for a cocaine-associated stimulus. *Psychopharmacology (Berl)* 126: 311–322.
- Whiteside SP, Lynam DR (2003). Understanding the role of impulsivity and externalizing psychopathology in alcohol abuse: application of the UPPS impulsive behavior scale. *Exp Clin Psychopharmacol* 11: 210–217.
- Wilens TE, Faraone SV, Biederman J, Gunawardene S (2003). Does stimulant therapy of attention-deficit/hyperactivity disorder beget later substance abuse? A meta-analytic review of the literature. *Pediatrics* 111: 179–185.
- Willuhn I, Wanat MJ, Clark JJ, Phillips PE (2010). Dopamine signaling in the nucleus accumbens of animals self-administering drugs of abuse. *Curr Top Behav Neurosci* 3: 29–71.
- Wilson AW, Neill JC, Costall B (1996). The 5-HT1A receptor agonist 8-OH-DPAT reduces ethanol intake and maintained behavior in female Sprague-Dawley rats. *Alcohol* 13: 407–413.

- Wilson TD, Dunn EW (2004). Self-knowledge: its limits, value, and potential for improvement. *Annu Rev Psychol* 55: 493–518.
- Winstanley CA, Dalley JW, Theobald DE, Robbins TW (2003a). Global 5-HT depletion attenuates the ability of amphetamine to decrease impulsive choice on a delay-discounting task in rats. *Psychopharmacology (Berl)* 170: 320–331.
- Winstanley CA, Chudasama Y, Dalley JW, Theobald DE, Glennon JC, Robbins TW (2003b). Intra-prefrontal 8-OH-DPAT and M100907 improve visuospatial attention and decrease impulsivity on the five-choice serial reaction time task in rats. *Psychopharmacology (Berl)* 167: 304–314.
- Winstanley CA, Dalley JW, Theobald DE, Robbins TW (2004). Fractionating impulsivity: contrasting effects of central 5-HT depletion on different measures of impulsive behavior. *Neuropsychopharmacology* 29: 1331–1343.
- Winstanley CA, Baunez C, Theobald DE, Robbins TW (2005a). Lesions to the subthalamic nucleus decrease impulsive choice but impair autoshaping in rats: the importance of the basal ganglia in Pavlovian conditioning and impulse control. *Eur J Neurosci* 21: 3107–3116.
- Winstanley CA, Theobald DE, Dalley JW, Robbins TW (2005b). Interactions between serotonin and dopamine in the control of impulsive choice in rats: therapeutic implications for impulse control disorders. *Neuropsychopharmacology* 30: 669–682.
- Winstanley CA, Eagle DM, Robbins TW (2006). Behavioral models of impulsivity in relation to ADHD: translation between clinical and preclinical studies. *Clin Psychol Rev* 26: 379–395.
- Winstanley CA, Bachtell RK, Theobald DE, Laali S, Green TA, Kumar A *et al.* (2009). Increased impulsivity during withdrawal from cocaine self-administration: role for DeltaFosB in the orbitofrontal cortex. *Cereb Cortex* 19: 435–444.
- Winstanley CA, Olsson P, Taylor JR, Jentsch JD (2010a). Insight into the relationship between impulsivity and substance abuse from studies using animal models. *Alcohol Clin Exp Res* 34: 1306–1318.
- Winstanley CA, Zeeb FD, Bedard A, Fu K, Lai B, Steele C *et al.* (2010b). Dopaminergic modulation of the orbitofrontal cortex affects attention, motivation and impulsive responding in rats performing the five-choice serial reaction time task. *Behav Brain Res* 210: 263–272.
- Wischhof L, Hollenstein KJ, Koch M (2011). Impulsive behaviour in rats induced by intracortical DOI infusions is antagonized by co-administration of an mGlu2/3 receptor agonist. *Behav Pharmacol* 22: 805–813.
- Wiskerke J, Stoop N, Schetters D, Schoffeleer AN, Pattij T (2011a). Cannabinoid CB1 receptor activation mediates the opposing effects of amphetamine on impulsive action and impulsive choice. *PLoS ONE* 6: e25856.
- Wiskerke J, Schetters D, van Es IE, van Mourik Y, den Hollander BR, Schoffeleer AN *et al.* (2011b). mu-Opioid receptors in the nucleus accumbens shell region mediate the effects of amphetamine on inhibitory control but not impulsive choice. *J Neurosci* 31: 262–272.
- Wiskerke J, van Mourik Y, Schetters D, Schoffeleer AN, Pattij T (2012). On the role of cannabinoid CB1- and mu-opioid receptors in motor impulsivity. *Front Pharmacol* 3: 108.
- de Wit H (2009). Impulsivity as a determinant and consequence of drug use: a review of underlying processes. *Addict Biol* 14: 22–31.
- Wogar MA, Bradshaw CM, Szabadi E (1993). Effect of lesions of the ascending 5-hydroxytryptaminergic pathways on choice between delayed reinforcers. *Psychopharmacology (Berl)* 111: 239–243.
- Wooters TE, Bardo MT (2011). Methylphenidate and fluphenazine, but not amphetamine, differentially affect impulsive choice in spontaneously hypertensive, Wistar-Kyoto and Sprague-Dawley rats. *Brain Res* 1396: 45–53.
- Wooters TE, Neugebauer NM, Rush CR, Bardo MT (2008). Methylphenidate enhances the abuse-related behavioral effects of nicotine in rats: intravenous self-administration, drug discrimination, and locomotor cross-sensitization. *Neuropsychopharmacology* 33: 1137–1148.
- Xi ZX, Gilbert J, Campos AC, Kline N, Ashby CR Jr, Hagan JJ *et al.* (2004). Blockade of mesolimbic dopamine D3 receptors inhibits stress-induced reinstatement of cocaine-seeking in rats. *Psychopharmacology (Berl)* 176: 57–65.
- Xi ZX, Kleitz HK, Deng X, Ladenheim B, Peng XQ, Li X *et al.* (2009). A single high dose of methamphetamine increases cocaine self-administration by depletion of striatal dopamine in rats. *Neuroscience* 161: 392–402.
- Xie X, Ramirez DR, Lasseter HC, Fuchs RA (2010). Effects of mGluR1 antagonism in the dorsal hippocampus on drug context-induced reinstatement of cocaine-seeking behavior in rats. *Psychopharmacology (Berl)* 208: 1–11.
- Xie X, Arguello AA, Reittinger AM, Wells AM, Fuchs RA (2012a). Role of nicotinic acetylcholine receptors in the effects of cocaine-paired contextual stimuli on impulsive decision making in rats. *Psychopharmacology (Berl)* 223: 271–279.
- Xie X, Lasseter HC, Ramirez DR, Ponds KL, Wells AM, Fuchs RA (2012b). Subregion-specific role of glutamate receptors in the nucleus accumbens on drug context-induced reinstatement of cocaine-seeking behavior in rats. *Addict Biol* 17: 287–299.
- Yee J, Famous KR, Hopkins TJ, McMullen MC, Pierce RC, Schmidt HD (2011). Muscarinic acetylcholine receptors in the nucleus accumbens core and shell contribute to cocaine priming-induced reinstatement of drug seeking. *Eur J Pharmacol* 650: 596–604.
- Yonghui L, Xigeng Z, Yunjing B, Xiaoyan Y, Nan S (2006). Opposite effects of MK-801 on the expression of food and morphine-induced conditioned place preference in rats. *J Psychopharmacol* 20: 40–46.
- Zeeb FD, Floresco SB, Winstanley CA (2010). Contributions of the orbitofrontal cortex to impulsive choice: interactions with basal levels of impulsivity, dopamine signalling, and reward-related cues. *Psychopharmacology (Berl)* 211: 87–98.
- Zernicke KA, Cantrell H, Finn PR, Lucas J (2010). The association between earlier age of first drink, disinhibited personality, and externalizing psychopathology in young adults. *Addict Behav* 35: 414–418.
- Zhang XY, Kosten TA (2005). Prazosin, an alpha-1 adrenergic antagonist, reduces cocaine-induced reinstatement of drug-seeking. *Biol Psychiatry* 57: 1202–1204.
- Zislin G, Desai TV, Prado M, Shah HP, Bruijnzeel AW (2007). Effects of the CRF receptor antagonist D-Phe CRF(12–41) and the alpha2-adrenergic receptor agonist clonidine on stress-induced reinstatement of nicotine-seeking behavior in rats. *Neuropharmacology* 53: 958–966.